GENETICS

A PERIODICAL RECORD OF INVESTIGATIONS BEARING
ON HEREDITY AND VARIATION

VOLUME 5 1920

PUBLISHED BY
THE PRINCETON UNIVERSITY PRESS
PRINCETON, NEW JERSEY
AND
WILLIAMS & WILKINS COMPANY
BALTIMORE

Second Printing 1965 / University of Texas Printing Division, Austin

GENETICS

A PERIODICAL RECORD OF INVESTIGATIONS BEARING ON HEREDITY AND VARIATION

EDITORIAL BOARD

GEORGE H. SHULL, Managing Editor
Princeton University

WILLIAM E. CASTLE
Harvard University

EDWARD M. EAST
Harvard University

EDWIN G. CONKLIN
Princeton University

ROLLINS A. EMERSON
Cornell University

CHARLES B. DAVENPORT
Carnegie Institution of Washington

HERBERT S. JENNINGS
Johns Hopkins University

BRADLEY M. DAVIS
University of Michigan

THOMAS H. MORGAN
Columbia University

RAYMOND PEARL
Johns Hopkins University

WILLIAMS AND WILKINS COMPANY
BALTIMORE
SANS TÂCHE

PRINCETON
UNIVERSITY
PRESS

TABLE OF CONTENTS

JANUARY NUMBER

(Issued May 17, 1920)

Portrait of JOSEPH GOTTLIEB KÖLREUTER ..... Frontispiece

Page

ALTENBURG, EDGAR, and MULLER, HERMANN J., The genetic basis of truncate wing,—an inconstant and modifiable character in Drosophila 1

WARREN, DON C., Spotting inheritance in *Drosophila busckii* Coq.... 60

MARCH NUMBER

(Issued August 10, 1920)

GOWEN, JOHN W., Studies in milk secretion, V. On the variations and correlations of milk secretion with age 111

KELLY, JAMES P., A genetical study of flower form and flower color in *Phlox Drummondii* 189

MAY NUMBER

(Issued September 9, 1920)

GOWEN, JOHN W., Studies in milk secretion. VI. On the variations and correlations of butter-fat percentage with age in Jersey cattle 249

DUNN, L. C., Linkage in mice and rats 325

DUNN, L. C., Independent genes in mice 344

JULY NUMBER

(Issued December 30, 1920)

ADKINSON, JUNE, The behavior of bronchial asthma as an inherited character 363

BLAKESLEE, A. F., A dwarf mutation in Portulaca, showing vegetative reversions 419

TABLE OF CONTENTS

(CONTINUED)

SEPTEMBER NUMBER

(Issued February 10, 1921)

Page

LAUGHLIN, HARRY H., Calculating ancestral influence in man: a mathematical measure of the facts of bisexual heredity... 435

SAFIR, SHELLEY R., Genetic and cytological examination of the phenomena of primary non-disjunction in *Drosophila melanogaster*..... 459

STURTEVANT, A. H., Genetic studies on *Drosophila simulans*. I. Introduction. Hybrids with *Drosophila melanogaster*..... 488

NOVEMBER NUMBER

(Issued March 4, 1921)

PAYNE, FERNANDUS, Selection for high and low bristle number in the mutant strain "reduced"..... 501

EAST, E. M., and JONES, D. F., Genetic studies on the protein content in maize..... 543

Index..... 611

CORRIGENDA

Page 18, line 10, for "wheras" read "whereas", and for "T'_3" read "T'_2".
Page 30, line 10, for "germ splasm" read "germ plasm".
Page 38, line 2 from bottom, for "lines 1 and 2" read "line 1".
Page 44, line 9 from bottom, "41" should be light-face like the denominator.
Page 60, line 7 of table of contents, for "Selection in line 500....64" read "Selection line 500 .... 66".
Page 60, line 8 of table of contents, for "67" read "69".
Page 60, line 9 of table of contents, for "68" read "70".
Page 60, line 10 of table of contents, for "70" read "72".
Page 60, line 11 of table of contents, for "71" read "73".
Page 60, line 12 of table of contents, for "73" read "75".
Page 60, line 13 of table of contents, for "74" read "76".
Page 60, lines 14 and 15 of table of contents for "75" read "77".
Page 180, line 15 of text, "10.648" should be raised to the position of an exponent of the quantity in parenthesis.
Page 190, line 20, insert semi-colon after the word "present".
Page 195, figure 4 is inverted.
Page 195, line 2 below the figure, for "*Pholox*" read "*Phlox*".
Page 211, line 9, for "magneta" read "magenta".
Page 215, last 3 lines of first column, for "$(AB)_{251.7}$" read "$(AB)_{250.20}$" for "$(AB)_{250.20}$" read $(BA)_{269.11}$"; and for "$(AB)_{269.11}$" read $(AB)_{251.7}$".
Page 217, line 9, for "anthocayanin" read "anthocyanin".
Page 222, line 6 from bottom, for "$(BA)_{269.1}$" read" $(BA)_{269.12}$".
Page 223, line 4 below table, for "5" read "25".
Page 228, line 11, insert comma after "D_e".
Page 237, line 7, for ".8 of" read "about".
Page 239, lines 1 and 2 below table, at right only, for "III a" read "III b".
Page 246, line 6 of paragraph 5b, for "*P*" read "*P* or *D*"
Page 247, line 6 of Literature cited, for "131 pp." read "154 pp."
Page 340, graph of chromosome, at extreme right only, for "*p*" read"*p'*".
Page 340, line 1 below graph, for "th" read "the".
Page 353, heading of table 5, third column from left, insert "and" before "Type A."

CORRIGENDA (CONTINUED)

Page 355, table, 6, between columns 1 and 2, insert a column as follows:

Agouti b.-e.-white *A W*
64 *76.5*

Page 355, column 3 of table 6, for "41" read "76," and for "*38.25*" read "*76.5*".

Page 355, column 4 of table 6, for "76" read "64".

Page 458, plate 2, a dotted line should connect P_5, No. 15 with P_4, No. 8.

Page 462, line 3, for "minature" read "miniature".

Page 466, line 6 from bottom, for "page 000" read "page 478".

Page 467, line 14 from bottom, for "minature" read "miniature".

Page 476, line 13 from bottom, for "othe frraction" read "other fraction".

Page 481, line 4 from bottom, for "minature" read "miniature".

Page 486, line 3 of paragraph 8, for "minature" read "miniature".

Page 558, line 7 from bottom, for "F" read "F_2".

Page 575, line 4 below table, for "20A–8–5–28" read "20A–8–5–22".

Dr. Rölmuster.

JOSEPH GOTTLIEB KÖLREUTER

(FRONTISPIECE)

This first great geneticist was born at Sulz-am-Neckar, Württemberg, Germany, April 27, 1733, and died at Karlsruhe, Baden, November 12, 1806. In 1748 he entered the UNIVERSITY OF TÜBINGEN, from which he received the degree of Doctor of Medicine in 1755, having spent one of the intervening years at the UNIVERSITY OF STRASSBURG. In 1756 he went to St. Petersburg where he had charge of the extensive collections of the IMPERIAL ACADEMY OF SCIENCE, and while there published some 20 notes and brief memoirs, chiefly on zoölogical subjects. In 1761 he returned to Germany as Professor of Natural History, first at Calw, and then in 1764 at Karlsruhe, where he was appointed Director of the Gardens of the Margrave KARL FRIEDRICH, a position which was rendered unhappy and finally untenable by the jealousy of the gardeners whom he superseded in authority. Although he ceased to be Director of the Gardens in 1769, he continued under the patronage of the Margrave throughout the rest of his life.

While at St. Petersburg, KÖLREUTER achieved success in the production of plant hybrids. His cross between *Nicotiana rustica* ♀ and *Nicotiana paniculata* ♂ was made in 1760, and 78 of the young hybrids were grown the same year. Twenty of these were brought to flower in-doors in March, 1761. With the exception of a single cross made by LINNAEUS (1757) between two species of Tragopogon, these were the first plant hybrids ever produced as the result of a fundamental genetical experiment, though at least one authentic hybrid was known to horticulture as early as 1719 (FAIRCHILD'S pink).

These hybrids were intermediate between their parents in all the more striking differential characters, and the pollen was scanty or wanting. Similar phenomena were found to be generally the characteristics of species hybrids, and while there are some notable exceptions, the occurrence of pollen sterility in hybrids has been recognized as an important index of the specific distinctness of the parents, and this test is known as "Kölreuter's method."

KÖLREUTER'S second Nicotiana cross (*Nicotiana transylvanica* × *N. glutinosa*), stands as the classic example of the excessive vigor often seen in hybrids, the phenomenon now known as "heterosis."

While most of KÖLREUTER'S hybrids were identical in reciprocal crosses, he secured notably dissimilar reciprocals in Digitalis; and in Verbascum he met and recognized the phenomenon of self-sterility. In one of his later papers he also distinguished between "constant" and "inconstant" hybrids, but evidently did not appreciate the importance of this distinction. J. BEHRENS, who has prepared the only comprehensive biography of KÖLREUTER,* credits him with 283 successful hybridization experiments, of which the greater number involved Mirabilis (82), Dianthus (54), Nicotiana (43), Verbascum (36), and the Malvaceae (20).

Besides his strictly genetical experiments, KÖLREUTER made careful studies of pollen, of whose form and structure he gave descriptions, but owing to the inade-

* BEHRENS, J., Joseph Gottlieb Koelreuter. Ein Karlsruher Botaniker des 18. Jahrhundert. Verhandl. d. naturwiss. Vereins in Karlsruhe **11**: 268-320. *1 portrait.* 1894.

quacy of the microscopes then available, he did not discover pollen-tubes, and so gained an erroneous conception of the nature of the fertilization process, which he thought to be a mingling of a female fluid secreted by the stigmas and a male fluid expressed from the pollen grains.

He also made numerous observations and experiments on nectar, and recognized and stated repeatedly and with emphasis the relation of insects to plant pollination. In this he anticipated by nearly 30 years the brilliant work of SPRENGEL (1793). SPRENGEL'S "Discovered secret of nature" was no secret to KÖLREUTER nor to those who had read the published accounts of his experiments.

Only four of KÖLREUTER'S papers are generally known, namely, his "Vorläufige Nachricht von einigen das Geschlecht der Pflanzen betreffenden Versuchen und Beobachtungen," and three "Fortsetzungen," which were published, respectively, in 1761, 1763, 1764 and 1766. These four papers were republished in 1893 in Ostwald's Klassiker. A note in the latter intimates that these were the only papers published by KÖLREUTER on his hybridization experiments, but this is erroneous, as he wrote between 1772 and the time of his death, 22 botanical and several zoölogical papers, mostly relating to hybridizations in various groups of plants. Nearly all of these were published in the Acta Academiae Scientiarum Imperialis Petropolitanae, St. Petersburg.

KÖLREUTER'S strict adherence to the experimental method, his carefully planned and executed experiments, his direct and lucid statements of results, and the clear logic of his inductions and deductions, were strictly modern and in striking contrast to the writings of most of his contemporaries.

The portrait here reproduced is the only one of KÖLREUTER generally known, and it has been repeatedly engraved. We have been unable to secure information as to the present location of the oil painting which forms the basis of all these various engravings, but in 1894, when it was first reproduced, it was in the possession of W. KÖLREUTER, a grandson of J. G. KÖLREUTER, Hornberg, Baden, and it is probably still in the family. A second oil painting which is practically a mirror image of this one, is included among the relics of C. F. GAERTNER which were deposited by GAERTNER'S daughter in the Botanischen Institut of the UNIVERSITY OF TÜBINGEN. It is not known whether this second portrait is an original or whether it was copied from the one already mentioned, but Professor E. LEHMANN to whom we are indebted for photographic copy of this potrait is inclined to believe that it is the original and the only *original* portrait of KÖLREUTER. He suggests that the facing of the engraved portrait to the right instead of the left was a result of the process of engraving. If this explanation is correct, certain other minor differences must be explained in the same manner.

The engraving used as the basis of our frontispiece was retouched and changed from an oval to an oblong setting by J. MARION SHULL, of the U. S. DEPARTMENT OF AGRICULTURE. The fund necessary for its reproduction was generously provided by Professor OAKES AMES, of HARVARD UNIVERSITY, well known for his numerous papers on the Orchidaceae.

THE GENETIC BASIS OF TRUNCATE WING,—AN INCONSTANT AND MODIFIABLE CHARACTER IN DROSOPHILA

EDGAR ALTENBURG
The Rice Institute, Houston, Texas
and
HERMANN J. MULLER
Columbia University, New York City

[Received August 4, 1919]

TABLE OF CONTENTS

The initial selection experiments and crosses 1
Analysis of the effects of the different chromosomes in truncate stock, on the truncate character 7
Tests of the constancy of the factors for truncate 20
a. Tests of sibs 20
b. The "pure-line" experiment 25
The cause of inconstancy of truncate stock 30
a. Proof of the viability and non-sterility of flies homozygous for the intensifiers 30
b. Experiments to test the possibility of securing homozygous T_2' 36
c. The reason for the non-appearance of homozygous T_2' 39
d. The cause of the infrequency of normals in the selected stock 40
e. The source of the normals which did occur in the selected stock 42
The locus of truncate 43
On the mode of origin of the truncate complex 46
The origination of truncate in other stocks 49
Applications of the methods 51
RESUMÉ 53
LITERATURE CITED 59

THE INITIAL SELECTION EXPERIMENTS AND CROSSES

One of the first mutants which MORGAN found in Drosophila was a fly with truncated wings. It appeared in 1910, in a stock having beaded wings, and the inheritance of truncate, like that of beaded, seemed from the first to be irreconcilable with Mendelian principles, or indeed with any theory of fixed and segregating factors.

The truncate and the beaded cases thus stood as a constant challenge to those workers who hoped for a completely rational explanation of in-

heritance in Drosophila. Beaded has been dealt with in another paper; the truncate case will be considered here.[1]

The original truncate male, when crossed by its normal sisters, gave, even in F_1, a motley collection of offspring, the great majority being normal, but truncates of various grades also appearing. In F_2 the results were varied, and some of the F_1 matings of truncate by truncate gave nearly half the F_2 flies having some degree of truncation. It was to be expected that in the next generation pure stocks could be established; nevertheless crosses of the F_2 failed to produce any higher proportion of truncate than before, and for a few generations it seemed as though selection might be ineffectual. Broods yielding a higher percentage were found after several months, however, and by carrying on the selection work with these lines for about a year, MORGAN finally was able to obtain a stock in which sometimes as much as 90 percent of the flies showed the mutant character. The intensity of the character had increased markedly at the same time, but was accompanied by a greatly diminished fertility of the flies.

At this point one of the authors took up the work, and in an effort to obtain pure stock the process of selection was continued for nearly three years longer. It seemed, however, as though the climax had already been reached, for it was found impossible entirely to eliminate the normal individuals from the stock, or to obtain races that consistently yielded more than about 90 percent of truncates of all grades. It would not be justifiable to publish the protocols of these experiments in detail as this sort of breeding work is necessarily of an uncritical nature, not adapted to give definite factorial solutions to genetic problems. Table 1, however, is given as a sample of this series of experiments; it shows the effects simply of 13 successive generations of the selection, carried on during the latter part of the period. It will be seen that the average proportion of truncates (all grades) shows no significant change, being 84 percent for the first four generations, 90 percent for the intermediate five, and 88 percent for the last four.

These results suggest that this final stock might really be pure genetically, and homozygous for truncate, and that the differences between the individuals were due only to external conditions which accentuated or inhibited the development of the somatic character. In that case the normals and the truncates of various grades should all give identical results

[1] A preliminary account of the truncate case was given by the authors in the book of MORGAN, STURTEVANT, MULLER, and BRIDGES (1915, pp. 191-194). Most of the crosses there reported were made in 1913.



FIGURE 1.—*Types of truncation.*

a. Very short truncate male (the blistering is a frequent characteristic of this grade).
b. Typical short truncate male of the selected stock.
c. Long truncate female—the type of "truncate" usual in outcrosses.
d. Intermediate-winged female.
e. "Slight intermediate" male.
f. Normal-winged male from wild-type stock.

when bred. Table 2, however, demonstrates the wide genetic difference between the normals and the truncates thrown by this stock, for whereas the truncates on the average had given 91 percent of truncates in this experiment, the normals yielded only 5 percent. Even the selected stock therefore is undergoing a continual genetic variation, with which the variation in the character is correlated, and so selection back and forth within this stock still has immediate and marked effect, within the 90 percent limit.

TABLE 1

"Main line" of truncates selected for 13 generations for homozygosis.

Number of generation	Truncates (all grades)	Normals	Average grade of truncation according to generation	Average grade according to groups of generations	Average percentage of truncates
1	*13	0	3.0	2.52	84
2	5	0	2.3		
	*19	4			
	16	8			
3	*11	0	2.7		
	15	1			
	12	3			
4	*36	8	2.5		
5	*18	2	2.7	2.70	90
6	23	3	2.5		
	8	2			
	*28	8			
7	*98	6	2.8		
	9	2			
	53	6			
8	*13	2	2.6		
9	*8	0	3.0		
10	30	3	2.6	2.64	88
	*19	4			
11	10	1	2.5		
	*29	7			
12	*79	10	2.7		
13	28	1	2.9		

* The asterisks denote those families from which the parents of the families recorded in the next generation were taken. It will be seen that, in some cases, as in generation 6, the family thus marked out is not the one that shows the highest percentage of truncates. Such a circumstance is due to the infertility of the "best-truncate" families; attempts were made in each generation to breed from the "best" families, as well as from the others, but the "best" lines frequently died out after one or several genera-

tions, the race then being continued from the families of those poorer lines which had been carried on simultaneously as a safeguard. The above table shows none of the unsuccessful side lines which were in this manner always being carried on, both from better and from poorer stocks, but it gives a record only of those families that happened to lie in the line that was carried through continuously.

As a typical example of an unsuccessful side line, it may be stated that in generation 6, referred to above, three cultures were made of the truncate flies in the "best family" (that which consisted of 23 truncates, 3 normals). The "best" of these three cultures gave, in generation 7, 78 truncates and 1 normal; two families in turn were started from this culture; of these, one produced nothing at all, and the other yielded only three infertile flies in generation 8. The "main line" thus reverted to the descendants of the third bottle of generation 6, which has therefore been marked with the asterisk.

In determining the average grade of truncation of the above counts, the class of truncate is assigned an average grade of 3 and the class of normals a grade of 0, each grade is then multiplied by the number of flies in the class having that grade, and the sum of these products is divided by the total number of flies.

TABLE 2

Genetic behavior of normals thrown by truncates of the selected stock.

Number of generation	Truncates	Normals
1	38	3
2	21	1
3	17	1
4	62	9
5	0	27
6	1	19
7	5	67

Lines 1 to 4 show the counts given by the truncate stock for the four generations preceding that in which normals were chosen; only those families are here recorded which are in direct line with the one (4) from which the normals were derived. Lines 5 to 7 show the ratios thrown by the normals during three successive generations in which they were selected as parents. Individual matings (separate pairs of flies) were not used in this experiment.

The results of crossing truncate to wild-type flies were extraordinarily indefinite and inconclusive. A few examples of the numerous crosses of this sort which were carried out are given in table 3. Truncate behaves nearly as a recessive in F_1 (see table 3(A)), although a small percentage of flies appeared with the character more or less perfectly developed. In F_2 a good 3:1 proportion was never secured, but, as may be seen from table 3(B), ratios occurred varying from 168 normals: 1 truncate, to less than 2 normals: 1 truncate, the "truncates" being of various grades. In other crosses of the same type F_2 families have also occurred in which no truncates were produced at all. When an attempt was made to recover truncate stock by breeding together the "extracted" truncates

TABLE 3

Crosses of truncate by wild stock.

(A.) F_1 (from wild female by truncate male)

Bottle No.	Character of parents used	Truncate ♀	Truncate ♂	Intermediate ♀	Intermediate ♂	Normal ♀	Normal ♂	Percentage of truncates (all grades)
1	Wild ♀, T ♂ from selected stock	5	1	4	2	162	182	3.36
2	Wild ♀, T ♂ from selected stock	5	0	1	3	129	127	3.4
3	Wild ♀, T ♂ from selected stock	0	1	1	0	89	83	1.15
4	Wild ♀, T ♂ from selected stock	4	1	1	3	214 (♀&♂)		4.03
5	Wild ♀, T ♂ from early unselected stock	0	0	0	0	672	676	0.0

(B.) F_2 (from $F_1 \times F_1$)

Bottle No.	Character of parents used	Truncate ♀	Truncate ♂	Intermediate ♀	Intermediate ♂	Normal ♀	Normal ♂	Percentage of truncates (all grades)
6	T ♀, normal ♂ from 3	22	9	3	4	103	68	18.2
7	" " " " " 4	15	11	7	1	32	21	39.2
8	I " " " " 3	28	15	20	14	173	120	20.8
9	N ♀, N ♂ both from 3	17	6	3	3	184	174	7.5
10	N ♀, N ♂ " " 5	0	2	0	0	208	128	.6
11	" " " " " " 5	12	20	3	5	168	125	12.0

(C.) F_3 (from $F_2 \times F_2$)

Bottle No.	Character of parents used	Truncate ♀	Truncate ♂	Intermediate ♀	Intermediate ♂	Normal ♀	Normal ♂	Percentage of truncates (all grades)
12	T ♀, T ♂ both from 6	85	49	14	13	54	42	62.6
13	" " " " " " 8	101	43	13	10	77	111	47.04
14	N ♀, N ♂ " " 8	37	22	23	22	158	172	23.0

(D.) F_4 (from $F_3 \times F_3$)

Bottle No.	Character of parents used	Truncate ♀	Truncate ♂	Intermediate ♀	Intermediate ♂	Normal ♀	Normal ♂	Percentage of truncates (all grades)
15	T ♀, T ♂ both from 12	132 (♀&♂)		11 (♀&♂)		44 (♀&♂)		76.5
16	" " " " " " "	79	58	43	42	1	8	63.0
17	" " " " " " "	32	17	1	0	112 (♀&♂)		30.9
18	N ♀, N ♂ " " "	35	31	4	5	50	73	37.8
19	" " " " " " "	0	0	0	0	464 (♀&♂)		0.0

of the F_2 families, it was found (table 3(C)) that truncate had been considerably "weakened," as only about 50 to 60 percent of truncates were now produced. Further selection, however, brought the extracted stock in later generations nearer to the value of the original selected stock. (An indication of this may be seen in table 3(D).) It was found that the percentage of truncates in F_2 and later generations of the cross tended to be higher if the P_1 truncate fly had been a female, than if the reciprocal P_1 cross had been made, and yet fair numbers of truncates of each sex were produced in both cases. All these results could be "explained" as due to varying "potencies," but this did not put the matter on a factorial basis.

Besides being crossed to wild-type, truncate was also crossed to various mutant stocks, and counts were made of the F_1, and of the F_2 derived from interbred F_1. Here the same indefinite results were obtained, in respect to the truncate wing character, as in the crosses with wild, but in addition there was also observable an apparently indefinite sort of linkage between truncate and the other character, which appeared, curiously enough, regardless of the linkage group to which the other character belonged. As examples of such experiments, the results of some of the crosses with jaunty (group II), and with pink (group III), are given in tables 4 and 5 respectively. It will be seen that in the F_2 of the cross with pink, although some degree of linkage was clearly indicated, yet a surprisingly large proportion of pink truncates were produced in certain bottles,—more than could be accounted for by weakness of linkage and by the amount of dominance shown by truncate in F_1.

ANALYSIS OF THE EFFECTS OF THE DIFFERENT CHROMOSOMES IN TRUNCATE STOCK ON THE TRUNCATE CHARACTER

It was thus necessary to discover, first, whether truncate depended on definite factors, lying in the chromosomes, and, secondly, if this proved to be true, whether the continual genetic variation occurring in the stock was due to fluctuation in these factors, or to heterozygosis that for some reason was forced to persist in the stock. As it was evident that at any rate the results did not represent simple monohybrid ratios, a method was required whereby the different factors, or sets of factors, that might be concerned in producing truncate could be dissociated from each other and studied separately. This object was achieved by making use, in a special way, of the linkage relations which had been worked out in Drosophila.

TABLE 4

Crosses of truncate by jaunty.

Generation of count	Bottle number	Character of parents ♀	Character of parents ♂	Straight-winged: Truncate ♀	Truncate ♂	Inter. ♀	Inter. ♂	Normal* ♀	Normal* ♂	Jaunty-winged (all normal*) ♀	Jaunty-winged (all normal*) ♂	Percentage T and I among straight
F_1	1	jaunty	T	1	0	16	2	120	141			6.8
F_1	2	T	jaunty	0	4	1	1	17	19			14.3
F_1	3	T	jaunty	0	4	9	6	66	38			15.4
F_2	4	N from 1	N from 1	1	2	1	1	58	63	8	9	4.0
F_2	5	N from 1	N from 1	5	9	2	0	91	79	15	17	8.6
F_2	6	I from 1	I from 1	21	10	8	3	19	43	14	7	40.4
F_2	7	N from 2	N from 2	4	0	2	0	29	30	11	4	9.2
F_2	8	N from 3	N from 3	13	4	9	1	72	113	7	7	12.7
Sum of F_2				44	25	22	5	269	328	55	44	13.8

* The word "normal," or "N," in this and the following tables, means normal in respect to truncation.

It may be recalled that in Drosophila all the hereditary factors are divisible into four groups, corresponding to, or rather contained in, the four pairs of chromosomes, and that in the male all the factors within any one particular chromosome are inseparable from one another during segregation (no crossing over), so that every offspring which inherits a single factor of that chromosome necessarily inherits also all the other factors which were in that particular chromosome. That is, all the factors lying within a given chromosome of a male have an identical distribution among his offspring. Thus it is only necessary to be able to observe the distribution, among the offspring, of a single easily recognizable pair of characters from the male parent, in order to learn the

TABLE 5

Crosses of truncate by pink.

Generation of count	Bottle No.	Character of parents ♀	Character of parents ♂	Red-eyed: Truncate ♀	Red-eyed: Truncate ♂	Red-eyed: Intermediate ♀	Red-eyed: Intermediate ♂	Red-eyed: Normal ♀	Red-eyed: Normal ♂	Red-eyed: Percent T+I among reds	Pink-eyed: Truncate ♂	Pink-eyed: Truncate ♀	Pink-eyed: Intermediate ♀	Pink-eyed: Intermediate ♂	Pink-eyed: Normal ♀	Pink-eyed: Normal ♂	Pink-eyed: Percent T+I among pinks
F_1	1	pink	T	4	2	9	2	184	227	4.0							
"	2	pink	T	16	10	9	2	66	86	19.6							
"	3	T	pink	1	3	3	6	57	66	9.6							
"	4	T	pink	2	6	10	9	88	101	12.6							
F_2	5	N from 1	N from 1	7	5	2	6	77	86	10.9	7	1	3	0	28	29	16.2
"	6	T from 1	N from 1	12	3	7	4	54	48	20.3	0	0	0	0	18	12	0
"	7	I from 1	N from 1	6	8	1	4	56	47	15.6	0	0	0	0	11	13	0
"	8	N from 2	N from 2	6	2	0	2	101	105	4.6	0	0	0	0	14	16	0
"	9	N from 3	N from 3	23	10	3	3	128	108	14.2	0	0	1	2	33	40	4.0
"	10	N from 4	N from 4	7	6	7	6	75	87	13.8	1	0	2	0	22	25	6.0
"	11	I from 4	I from 4	23	13	10	11	38	41	41.9	0	0	0	0	13	15	0
"	12	I from 4	I from 4	28	23	11	12	39	57	43.5	5	3	5	5	20	32	25.7
"	13	I from 4	I from 4	40	27	12	11	61	82	38.5	1	0	4	2	11	13	22.6
"	14	I from 4	I from 4	14	12	7	4	18	23	47.5	1	0	0	1	5	6	15.4
Sum of F_2				166	109	60	63	627	684	23.2	15	4	15	10	175	201	10.4

distribution of all the factors lying in the corresponding pair of chromosomes. This will tell nothing, however, about the distribution of the factors in the other three pairs of chromosomes, since the different pairs of chromosomes are distributed independently of one another to the offspring, and so there is opportunity for factors in different pairs of chromosomes, which may in previous generations have accompanied each other, to become dissociated from one another—i.e., transmitted to different offspring. Hence in order to know what was the distribution of all four pairs of chromosomes of a male parent among his offspring, it is necessary to have under observation the inheritance of four pairs of characters—one for each of the chromosome-pairs; each of these characters, then, will serve for the identification of an entire chromosome or group of factors in the offspring, and may be called an "identifying character." So, by this method, it is possible to determine the distribution, among the offspring, of the entire constellation of factors which a male parent contained. The offspring will contain different combinations of his factors (assuming he was heterozygous), but in each case it will be known just what combination is present. The somatic effects of these different known combinations of factors on the offspring may then be studied, and so it may be determined just what rôle each of the groups of factors plays in the production of any particular character that is under investigation. In this manner it was possible to ascertain what effect on the development of truncate wing each chromosome of the truncate fly produced.

It should be noted, however, that the simplicity of such an experiment is impaired when the character under investigation (e.g. truncate) is recessive, for then the factors for it must be introduced from the female parent as well as from the male, in order that the character may be observable in the offspring. The disadvantage of thus introducing the factors from the female parent is that in the female the factors for the character under investigation are liable to cross over from the identifying factors of the groups to which they belong, and it may therefore be impossible to tell, by means of one identifying factor in each chromosome, exactly what combination of factors for the character the offspring studied have received.[2] It would accordingly be desirable to have the character dominant, in order that in the investigation the factors for it

[2] If, however, an identifying factor at each end of the chromosomes is used, it may be assumed with a fair degree of probability that the offspring receiving both identifying factors have also received those factors for the character studied which lay between them.

may be introduced from the male parent alone. In the case of truncate, this mode of procedure was rendered possible by a special circumstance.

Ordinarily, truncate behaves most nearly like a recessive character, when truncates are crossed to wild-type flies, although a very small percentage of F_1 showing some degree of truncation are almost always produced. Examples of such crosses have been shown in table 3. Here the flies which have wings at all truncated have been classified, for greater accuracy, into two divisions—typical "truncates," and "intermediates," and it will be seen that the number in the two classes is approximately equal. The total proportion of truncates—including both grades—is 3.1 percent (in the crosses of truncate of the selected stock). When truncate was crossed to various other types of flies somewhat similar results were usually obtained, but when the cross was tried of truncate from the selected stock by flies with black body color, the outcome proved to be conspicuously different, as shown in table 6. Here it will be seen that there are distinctly more typical truncates than intermediates, and the proportion of truncates of all grades appearing in this cross is also very much higher, being in fact 24.9 percent (in the cross of truncate males by black females), as compared with the previous 3.1 percent (when truncate males were crossed to wild-type females). The truncation of the two series of F_1 flies may be compared also by obtaining a figure which may be called the "average truncation" of each group of individuals. The typical truncates may be assigned a

TABLE 6

Crosses of truncates by black.

(A) F_1 from black ♀♀ × truncate ♂♂

Bottle No.	Truncate		Intermediate		Normal		Percentage of T and I
	♀	♂	♀	♂	♀	♂	
1	35	0	11	13	90	116	22.3
2	57	0	25	18	113	162	26.6
3	55	1	22	12	121	149	25.0
Sum	147	1	58	43	324	427	24.9

Average truncation among ♀♀ 1.33, among ♂♂ .19, among all .79.

(B) F_1 from truncate ♀♀ by black ♂♂

Truncate		Intermediate		Normal		Percentage of T and I
14	11	7	0	25	12	46.4

Average truncation among ♀♀, 1.52; among ♂♂, 1.91; among all, 1.65.

grade or value of truncation of 4, in this calculation, the intermediates a value of 2, and the normals 0. Multiplying each of these values by the number of individuals having the value, adding, and then dividing by the total number of individuals, it is found that the average grade of truncation of the F_1 from black females by truncate males is .79, whereas the average grade of the F_1 in the cross of selected truncate with normal is .10. This relatively high grade of truncation for the F_1 flies has been obtained consistently, in numerous and varied crosses of truncate with flies of black body color. Moreover, although there is still a considerable proportion of normals present among the F_1 even in these crosses, it will be shown later that "high grade" truncate is really almost always dominant in individuals heterozygous for black. These results with black were in themselves noteworthy, for here there seemed to be a type of interaction of factors hitherto not described, wherein two non-allelomorphic factors, both ordinarily recessive, re-enforced each other so as to produce a visible effect in the individuals heterozygous for both, just as happens in crosses of multiple allelomorphs. It should be added that the black character, unlike the truncate character, was not affected by this interaction, and remained recessive. The facts were more important, however, in their bearing on the truncate case, for they disclosed a means whereby truncate might be used as a dominant character, in the desired analysis.

In order to simplify the account of the analysis, crosses are here reported in which only the three large groups of factors were followed,—the fourth chromosome, which was too small to be likely to contain factors for truncate, was followed, however, in other experiments, and proved not to be concerned in the case. The sexes were used as the pair of "identifying characters" for the first chromosome; black body color, *b*, versus *B*, the normal gray, for the second chromosome; and pink eye-color, *p*, versus *P*, the normal red, for the third chromosome. The procedure was as follows: A truncate female with normal body and eye colors was mated to a normal-winged male with black body and pink eyes. The F_1 males, the distribution of whose factors among the F_2 flies was to be followed, had the following chromosome and factor composition, indicating the chromosomes by numerals and representing those derived from the truncate parent with asterisks, and the homologous chromosomes, derived from the black pink parent, without asterisks:

$$\frac{\text{I}^*X}{\text{I}\ Y}\ \frac{\text{II}^*B}{\text{II}\ b}\ \frac{\text{III}^*P}{\text{III}\ p}$$

Inasmuch as these flies are heterozygous for black, it was to be expected that truncate would behave as a dominant, at least partially, and the count designated as "cross 1" in table 7 confirms this expectation, for 33.3 percent of the flies had somewhat truncated wings. It is not in this generation, however, but in the next, that the desired analytical results appear.

One of these F_1 flies (a male) was back-crossed to homozygous black pink females of the composition

$$\frac{IX}{IX}\ \frac{IIb}{IIb}\ \frac{IIIp}{IIIp}$$

in order to obtain offspring (F_2) having various known combinations of the chromosomes derived from truncate stock. As all these offspring (F_2) alike received the non-dominant combination *IX IIb IIIp* from their homozygous mother, the circumstance of receiving I**X* or *IY*, II**B* or *IIb*, and III**P* or *IIIp* from their heterozygous father determined whether their characters were female or male, gray or black, and red or pink, respectively. The visible characters accordingly revealed just what chromosomes of the truncate stock these flies had or had not received. Thus, the female gray red flies must have received from their father all of his truncate-derived chromosomes,—I**X*, II**B*, and III**P*, —while the male black pink flies could not have received any of these chromosomes, but must have gotten the non-truncate combination *IY IIb IIIp* instead. Corresponding with these facts, we find that the former flies (gray red females) are very largely truncate in wing shape, whereas all the latter (black pink males) have normal wings. These are the two extreme combinations; all the others received from the heterozygous father certain chromosomes derived from the truncate stock and others derived from the normal-winged (black pink) stock; a study of the amount of truncation shown by each of the different combinations may then reveal the rôle played by each chromosome in the production of the truncate character.

The results of the cross are designated as "cross 3" in table 7. There are eight possible combinations of the three pairs of identifying characters, and it will be seen that all of these combinations, or classes, appeared with about equal frequency, just as expected. In the case of each class the flies were further subdivided into four different grades, according to their wing shape. These grades (normal, "slight" intermediate, intermediate, and "truncate") may be given average values of 0, 1, 2.5, and 4, respectively, to bring them into proper relationship with

the three grades previously used. The fact most obvious on examining the records of the different classes, with regard to their amount of truncation, is that all of the black flies have normal wings; i.e., none of the flies in any of the last four combinations, which failed to receive II*, shows any indication of the truncate character; on the other hand, some at least of the flies in each of the first four, gray classes—the classes that did receive II*—had wings with some degree of truncation. This proves that there is a factor (or factors) for truncate in II*, i.e., in the second chromosome. This factor (factors?) in heterozygous condition is sufficient "by itself" to cause truncation, even if none of the other chromosomes in the fly are derived from truncate stock, but if this factor is not present, truncate cannot appear (at least, not as a dominant), even if one of the first and also one of the third chromosomes in the fly are derived from truncate stock. The factor for truncate in chromosome II is therefore the "chief" factor for truncate—being both necessary, and self-sufficient,[3] for the appearance of truncate (as a dominant). It may be designated as T_2'; its normal recessive allelomorph would then be t_2', and in our notation t_2' may ordinarily be understood to exist in any formula of the second chromosome in which T_2' is not represented.

T_2' is not the only factor for truncate whose existence is disclosed by this table. Comparison of the four classes which received T_2' (the gray classes) shows that truncate appeared in them with very different frequencies and intensities. It is evident that the red-eyed gray classes—those which contain III*P (the first two classes of the table)—contain more and better truncates than the corresponding pink-eyed gray classes —those having IIIp (the third and fourth classes of the table). The chromosome III* that came originally from truncate stock therefore contains a factor (or factors) for truncate also. This may be called T_3'; like T_2' it is dominant, inasmuch as it may produce an effect when in heterozygous condition. Unlike T_2' however, it may be called a "modifier" or "intensifier" rather than a chief factor, for it can produce an effect on the wing only in the gray classes, which already contain II*B (T_2').

It was then shown that even when an opportunity is given for it to exist in homozygous condition, T_3', without T_2', is unable to cause truncation. This was done by crossing the males and females from the black-bodied, red-eyed classes, to each other; the result is given in table 7, cross 5. The parent flies must all lack T_2' but are heterozygous for T_3'.

[3] Within the limits of the experiment.

Among the offspring, therefore, one-fourth should be homozygous for T_3', yet none of the flies had truncate wings. Three had a very slight unevenness of margin that approached the condition seen in slight intermediates, but this did not seem to be transmitted to their offspring. T_3', then, has practically no effect on the wing except in the presence of T_2'. Moreover, T_3' is not really necessary for the production of truncation, since some of the pink flies of the main experiment—flies that carried T_2' but could not have received T_3'—are truncate, although of course truncation in them is not so frequent nor so intense as when T_3' is present. T_3' thus fulfills the definition of an intensifier.

In a manner similar to that in which T_3' was deduced, we may also infer the existence of at least one intensifier, T_1', in the first chromosome (I*X) derived from the truncate stock. For the tables show that the female classes—i.e., the flies with I*X—contain a much larger propor-

TABLE 7

Crosses to determine the effects of the different chromosomes of selected truncate stock.

(A) Initial crosses of truncate by black pink

Generation of count	Number of cross	Nature of parents ♀	Nature of parents ♂	Gray red T	Gray red I	Gray red S	Gray red N	Gray pink T	Gray pink I	Gray pink S	Gray pink N	Black red T	Black red I	Black red S	Black red N	Black pink T	Black pink I	Black pink S	Black pink N
F_1	1	1 T from stock	Several bp from stock	3	8		18												
				4	7		26												
"	2	Several bp from stock	3 T from stock	13	10		39												
				1	12		56												

(B) Back-crosses to separate the effects of the chromosomes

F_2	3	Several bp from stock	1 T from 1	57	23	0	7	2	8	6	64	0	0	0	74	0	0	0	82
				2	17	12	47	1	4	4	62	0	0	0	67	0	0	0	73
"	4	1 T from 1	Several bp from stock	2	1	1	6	1	3	1	4	0	2	0	10	0	0	0	9
				0	1	1	2	0	5	1	4	0	1	0	4	0	0	2	2

TABLE 7 (continued)

(C) Crosses of derivatives of the separation cross

Generation of count	Number of cross	Nature of parents ♀	Nature of parents ♂	Gray red T I S N	Gray pink T I S N	Black red T I S N	Black pink T I S N
F_3	5	Several black red N from 3	Several black red N from 3			0 0 0 121 0 0 0 127	0 0 0 36 0 0 0 38
"	6	4 gray pink (2S&2N) from 3	1 black red N from 3	6 5 11 43 4 2 3 60	0 0 0 42 0 0 1 59	1 1 1 55 1 1 1 73	0 0 0 45 0 0 1 44
"	7	4 black red N from 3	1 gray pink S from 3	5 11 14 22 2 8 11 31	0 3 1 39 1 2 8 47	0 0 0 49 0 0 0 46	0 0 0 47 0 0 0 49
"	8	4 black pink N from 3	1 gray red S from 3	8 14 11 9 5 6 8 14	0 1 0 37 1 3 1 30	0 0 0 54 0 0 0 17	0 0 0 29 0 0 0 24
"	9	2 gray red I from 3	Several bp from stock	3 5 11 48 5 12 9 43	0 4 9 45 0 3 14 55	4 1 2 52 3 4 2 76	0 3 5 55 0 1 4 44

Females are recorded on the upper line, males on the lower, in the case of each cross. T = truncate, I = intermediate, S = "slight-intermediate," N = normal-winged; in section (A), both I and S have been grouped under I.

tion of truncates, and are more intensely truncated, than the corresponding male classes, provided of course that II**B* (T_2') is present.

It might be objected here, however, that in the case of the sex chromosome the difference between the truncation of the males and females may not be due to an intensifier present in I**X* and absent in I*X*, but that it may be due rather to sex difference, depending on the XX condition of the female and the XY condition of the male. This possibility was tested out by making reciprocal crosses of truncate to non-

truncate flies. As will be seen from tables 4, 5, 6, and 7, the male offspring are distinctly more truncated (in comparison with their heterozygous sisters), when they have truncate mothers than when they have truncate fathers. In table 6, for instance, there was an average grade of truncation of .19 among the male offspring of truncate males, whereas the male offspring of the truncate female had a grade of 1.91—ten times as high. The daughters were not very different in the two crosses, being 1.33 in the first, and 1.52 in the second. The sons must therefore be able to receive a factor, or factors, for truncate from their mother, which they cannot receive from their father,—that is, there are one or more dominant factors for truncate that obey the scheme of sex-linked inheritance. There must consequently be one or more intensifiers, T_1', in chromosome I*; the symbol of the normal allelomorph, t_1', will ordinarily not be represented, but its presence may be taken for granted in all X chromosomes in which T_1' is not shown. T_1', like T_3', cannot cause truncation, even in homozygous condition, unless T_2' is present.

In the crosses in which the female parent was truncate the male offspring were nearly always somewhat more truncated than the female, because the males are haploid for sex-linked factors and therefore appear like homozygotes, whereas the females contain factors from both parents and show some effect of the normal allelomorph received from the father. T_1', in other words, is not a completely dominant factor.

Among flies which are homozygous in regard to the sex-linked factors, on the other hand, no matter whether they are pure for the intensifier, or for its normal allelomorph, the males are on the average less truncated than the females. This effect is especially evident where the normal allelomorph of T_1' is present and is clearly shown in tables 9 and 10 (to be discussed later in another connection), wherein the males and females are both lacking in T_1' throughout, and are like each other so far as all non-sex-linked factors are concerned. In these crosses the female parents were non-truncated and the male parents were heterozygous for T_2' and T_3', but lacked T_1' (having resulted from a cross of truncate male by non-truncate female). In table 9 (A) and (B), for example, it is easily observable that there is a great preponderance of females among the truncate and "good-intermediate" offspring; the intermediate class has about equal numbers of females and males, and the slight-intermediate and normal classes contain almost exclusively males. The grade of truncation among the 914 females in these two tables averages 3.2, whereas that among the 864 males averages 1.6. In table 10, the average grade of truncation in the 535 females is 3.1, that in the 517 males is 1.3.

We may say, then, that the character truncate is partially sex-limited, or rather, *sex-influenced,* its development being favored more in the female sex than in the male. Or, to put the matter in factorial terms, the sex-factor itself, or some factor normally lying in the sex chromosome, intensifies the development of truncate, to a degree depending upon whether this factor is present in one "dose" or two, just as is the case with sex itself. This fact has nothing to do with the existence of a mutant intensifier, T_1', in the same chromosome.

After truncate had thus been analyzed into its chromosomal components, it was demonstrated that it could also be resynthesized by breeding together individuals of non-truncated or imperfectly truncated classes (gray pink females, black red males, etc.),—classes which, according to the analysis, should contain "complementary" components. Thus, the gray pink females should contain T_1' and T_2', and the black red males just T_3'. By breeding them together, then, as good truncates should be obtained as were found among the gray reds, in fact, the count should be rather similar to that obtained from the heterozygous gray red females by black pink males, except that here there is a chance for crossing over between black and truncate. The count, designated as cross 6, table 7, is quite according to this expectation. It will be noted here that the male parent had been normal-winged, and the females were only slight intermediates or normal (not possessing T_3'); the knowledge of what results would be produced with respect to truncate was thus entirely derived from the identifying factors. Another, somewhat similar cross, was of black red (long) females by a gray pink (slight-intermediate) male. Here the male has T_1' and T_2', and the female T_3'. The count is shown on line 7 of table 7, and again shows the high-grade gray red truncates. Black pink females derived from the main experiment, and therefore containing T_1', were likewise crossed to one of their complementary brothers, a gray red male ($T_2'T_3'$). The count, on line 8 of the same table, shows typical truncates, as before.

These results are all in striking contrast to those of other crosses, in which the parental classes (derived from the same cross as those above) did not supply complementary elements; compare, for example, the case where black reds were crossed to each other (table 7, cross 5), with the crosses above. Crosses involving still other combinations of these classes will be presented later. In each case, the results conformed to the expectation based on the analytical findings.

Let us now return to the three crosses of "complementaries," for we shall find that closer scrutiny of them will give us some further informa-

tion regarding the nature of the accessory factors for truncate. Crosses number six and eight, besides illustrating the reconstruction of truncate, provide an indication of whether the component designated as T_2' consists of a single factor, or a group of factors for truncate lying in the second chromosome. For, if T_2' is of multiple composition, a distinctly lower grade of truncate would be produced in cross six than in cross eight, since in the former cross T_2' is supplied to the offspring by their heterozygous mother, in which the process of crossing over would allow these supposed constituent factors of T_2' to become separated and scattered among different offspring, wheras in the latter cross T_3' comes from the father, and so its various factors would have to stay together (no crossing over), and re-enforce each other in the offspring, thus producing truncates of a higher grade than the average of the first cross. On the other hand, if T_2' is just a single factor, the truncates in the two crosses will of course be of the same grade. A test for T_2' alone is involved here, as the two crosses were identical (not reciprocal) so far as the chromosomes other than II were concerned, the cross six being $\frac{T_1'X}{X}\ \frac{T_2'B}{b}\ \frac{p}{p}$ female by $\frac{X}{Y}\ \frac{b}{b}\ \frac{T_3'P}{p}$ male, and the cross eight being $\frac{T_1'X}{X}\ \frac{b}{b}\ \frac{p}{p}$ female by $\frac{X}{Y}\ \frac{T_2'B}{b}\ \frac{T_3'P}{p}$ male. When the averages were calculated, it was found that the average truncation of the red truncated offspring (including all grades showing truncation) in cross six was 2.34 and that the average in cross eight was almost exactly the same, 2.33. Blacks and grays are both included in the average, since, on account of crossing over between black and T_2' in the third cross, the truncates are distributed among both black and gray classes. Taking all the truncated offspring,—black and gray, red and pink,—it is 2.26 in cross six and 2.34 in cross eight, a difference that is scarcely significant. These results then, indicate that T_2' is a single factor.

In a similar manner, comparison of crosses seven and eight give data regarding the composition of T_3', for in cross seven there is a chance of crossing over between the possible constituents of T_3', since T_3' is supplied by the female, whereas in cross eight T_3' comes from the male. while the crosses are identical so far as the other chromosomes are concerned. The grade of truncation of the red flies in cross seven was 1.97, and of all flies in that cross, 1.91. These rather low figures incomparison with values 2.33 and 2.34, that we have seen were obtained in cross eight, thus indicate that T_3' consists of more than one intensifying factor in the third chromosome. Other evidence for this was obtained in

later crosses, when it was found possible to obtain, by crossing over, stocks of pink truncate which there was reason to believe contained an intensifier in the third chromosome, that was not as marked in its effect as the original T_3'.

From the various experiments reported above, it is evident that truncate is a character that is particularly susceptible to modification by various factors, such as T_1', T_2', T_3', and "sex" (in the sense previously explained). Even the symbols T_1', T_2', T_3', moreover, probably do not represent separately all the mutant factors for truncate that exist in truncate stock, since, as has just been shown, it is likely that T_3' at least is composite. And, in addition to such specific factors for truncate, and sex, it should be remembered that there are mutant factors concerned with the development of other characters, which influence truncate at the same time. Thus, we have seen that the factor for black body-color must be regarded as an intensifier of truncate, as truly as T_1' or T_3'. There is evidence that a number of other previously known factors behave somewhat similarly. For instance, when a truncate male was placed in a bottle containing at the same time both a virgin bar-eyed female and two virgin round-eyed females (all non-truncate), the bar-eyed offspring that hatched in this bottle (bar is a dominant) were distinctly more truncated than the round-eyed offspring. As the environments of the two sets of offspring must have been exactly the same, the factor for bar, or some factor in the bar stock, must be a dominant intensifier of truncate. This result is shown in table 8. The average truncation of the bar offspring was 1.44, that of the rounds was 0.[4]

TABLE 8

Cross of a single truncate male (selected stock) by various females in a single bottle. One female parent bar (dominant, sex-linked), one vermilion miniature (recessive, sex-linked), one wild type; all three normal-winged. Count of F_1.

Females								Males											
Normal-eyed				Bar-eyed				Normal-eyed				Bar-eyed				Vermilion miniature			
T	I	S	N	T	I	S	N	T	I	S	N	T	I	S	N	T	I	S	N
0	0	0	73	24	3	2	28	0	0	0	65	15	4	2	45	0	0	0	20

[4] It may be added that in this experiment one of the round-eyed females had been a vermilion-eyed miniature-winged fly (*v m*), and this female left offspring as well as the other female, as shown by the presence of some *v m* sons (*v* and *m* are sex-linked), yet these two mutant factors, unlike bar, failed to intensify the truncate character.

A somewhat similar experiment was performed with the dominant factor for star eyes (star is in chromosome II). Heterozygous star females were crossed to truncate males (of a rather low-grade pink truncate stock containing that part of T_3' which had crossed over with pink). The count of non-star offspring was 100 normal males, 80 normal females, no truncates of any grade; the count of stars was: males, 69 normal-winged, 27 slight intermediate, 12 intermediate, 1 truncate; females, 103 normals, 36 slight intermediate, 25 intermediate, 16 truncate.

TESTS OF THE CONSTANCY OF THE FACTORS FOR TRUNCATE

a. *Tests of sibs*

Although it had thus been established that truncate is certainly a complex case of multiple factors, nevertheless this alone did not seem to explain satisfactorily the final failure of the persistent effort that had been made to secure homozygous stock. For, in an animal in which all the factors are linked into only four groups, it should not have taken so very many generations to secure complete homozygosis, no matter how many factors had been in heterozygous condition at the start. It therefore seemed important next to obtain a method of investigation in which the effects on truncate due to the segregation of the multiple heterozygous factors that had been found should be recognizable, and could be eliminated from consideration, so that it could be determined whether or not there was any residuum of genetic variability in this character, due to actual fluctuations in the genes themselves. It seemed possible to carry out this mode of attack by a continuation of the previous main experiment, reported on pages 11 to 15 (table 7(B)), wherein the various factors had been followed out according to their chromosomes. The "residuum of variation," if there were any, should appear among the different males or females contained within any one of the separate eye-color body-color classes shown in table 7, cross 3.

The gray red males, for example, should, according to the accepted scheme of inheritance in Drosophila, all have contained identical factors for truncate, for all alike had received, without crossing over, corresponding truncate-bearing second and third chromosomes from the same heterozygous male parent. Their composition was $\frac{Y}{X}\ \frac{T_2'\ B}{b}\ \frac{T_3'\ P}{p}$ as already noted. Yet, as we have seen, some of these males appeared truncate, others intermediate, and others normal. If, then, these differences between them were really genetic, the factors for truncate, having

been transmitted similarly to all of them, must have been subject to intrinsic variation. In order to determine whether the differences were genetic, the gray red males of various wing shapes needed merely to be bred to black pink females from stock; counts of the number of truncates of different grades among the offspring of the separate males would then reveal whether their character differences had a genetic basis.

The advantages of using the gray red males rather than the gray red females for the testing are threefold:—(1) a larger number of offspring can be obtained from individual males than from individual females; (2) when males are tested, since there is no crossing over between the factors for truncate and the "identifying factors," gray and red, the offspring which are gray and red like their parents furnish an ideal test of the truncate which was present in the original gray red parents, and only this class of offspring need be examined for this purpose; if female flies are tested, on the other hand, their factors for truncate become scattered more or less indiscriminately amongst all the different classes of offspring, because of crossing over. (Compare the distribution of truncate among the offspring recorded in table 7, cross 9, representing the tests of heterozygous gray red females, with the distribution among the offspring of heterozygous gray red males, shown in table 7, cross 8); (3) when males are tested, these gray red offspring, being descended substantially without recombination from their gray red fathers, may be used again for similar tests in the next generation.

Tests were therefore made of gray red males derived from crosses similar to that shown in table 7, cross 3. The results of these tests are shown in table 9, (A) and (B), the count from each male being shown on a separate line. Only those offspring of the tested males which, like the latter, are gray red, are recorded, for the reason explained in (2) of the paragraph above; the gray pinks, etc., appeared as expected, however. Table 9 (A) represents tests of males which we may call group A; these all derived their truncate-bearing chromosomes from the same heterozygous $\frac{Y}{X}\ \frac{T'_2B}{b}\ \frac{T'_3P}{p}$ father; they should therefore be genetically identical in truncation, if factors are constant. Table 9 (B)—group B—represents tests of males derived from a different father from that of group A; they may therefore be different from group A genetically. They should, however, be like one another, as they too are brothers to each other, and were derived from a single $\frac{Y}{X}\ \frac{T_2'B}{b}\ \frac{T_3'P}{p}$ father.

It will be seen, on comparing the results of the tests of the different brothers, either in group A or group B, that although there was a certain amount of "chance" variation in the proportion of truncates produced by the different males, nevertheless this variation was entirely uncorrelated with the grades of truncation in the wings of the male parents themselves. Thus, in group A, the four truncate-appearing males produced families with the following average amounts of truncation: 2.0, 2.2, 2.4, 2.6; the final average for the offspring of the truncate males therefore is *2.3*+. In the same group (A), the 3 "good intermediates" produced averages of 1.8, 2.5, and 2.8, respectively, which gives a final average of *2.5*+; the intermediates produced 2.15, 2.25, 2.4, 2.4, final average *2.3*; the 2 very slight intermediates and one normal gave 2.15, 2.47, 2.5, final average *2.4*—. In group B a similar situation exists, the families of the 2 truncates and 1 good intermediate together averaging 2.5—, those of the 3 intermediates also averaging 2.5—, and those of the 2 slight intermediates and 2 normals together averaging 2.7. It is here evident that the variation between the families is entirely an individual matter, not connected with the class of wing shape to which the parent belongs. For there is much greater variation between the families within each parental class than between the average values for the different classes. In no case do the latter values (within either group A or B) differ by as much as 0.3 grade, in spite of the fact that the parents themselves, of these various classes, present average differences ranging from 1.0 grade (in the minimum cases) to 4.0 grades (in the maximum). Moreover, those slight differences which do exist between the offspring averages have absolutely no tendency to be of the same sign as the parental differences. It must therefore be concluded that, so far as these tests can show, the gray red brothers which were tested had been alike genetically; the relatively slight deviations in truncation, between their families of offspring, were not of genetic origin, since they were in no way connected with the variations in truncation between the brothers themselves.

Besides the above tests of males, similar but less extensive tests of sister gray red females were also carried on, which had a composition similar to that of the males, but contained in addition heterozygous T_1'. These tests yielded results identical in kind with those of the males, except that the counts were smaller, and less critical in nature, as explained on p. 21. In these tests too there was a conspicuous lack of correlation

TABLE 9

Tests of brother gray red males ($\frac{Y}{X}\ \frac{T_1' B}{b}\ \frac{T_2' P}{p}$) *by black pink females.*

All tested males within a single group (A or B) are derived from a single father of same type as themselves by black pink mother. Only their gray red offspring are recorded; the numbers not in parenthesis give the total males and females, those in parenthesis give the females.

(A) (Group A)

Grade of truncation of tested male parent	Average grade of truncation of offspring	Truncate	Good intermediate	Intermediate	Slight intermediate	Normal
4 (truncate)...	2.0	20(20)	9(9)	32(13)	41(0)	2(0)
4 (truncate)...	2.2	22(21)	11(9)	22(9)	32(1)	1(0)
4 (truncate)...	2.4	12(12)	8(8)	6(4)	15(0)	0(0)
4 (truncate)...	2.6	51(49)	24(16)	56(12)	33(0)	0(0)
Total of truncates (4)....	2.34	105(102)	52(42)	116(38)	121(1)	3(0)
3 (good intermediate	1.8	5(5)	6(6)	17(11)	20(1)	2(0)
3 (good intermediate	2.5	17(15)	10(8)	18(4)	16(0)	0(0)
3 (good intermediate	2.8	52(45)	26(17)	59(9)	11(0)	0(0)
Total of good intermediate(3)	2.53	74(65)	42(31)	94(23)	47(1)	2(0)
2(intermediate)	2.15	14(14)	6(5)	24(14)	22(0)	1(0)
2(intermediate)	2.25	20(15)	5(4)	16(12)	28(0)	0(0)
2(intermediate)	2.4	20(20)	11(10)	34(17)	17(1)	0(0)
2(intermediate)	2.4	25(24)	11(8)	17(9)	28(0)	2(0)
Total intermediates (2)....	2.30	79(73)	33(27)	91(52)	95(1)	3(0)
0.5 (very slight intermediate)	2.15	14(14)	7(7)	25(11)	19(1)	3(0)
0.5 (*very* slight intermediate)	2.5	29(22)	18(15)	50(25)	20(0)	0(0)
0.0 (normal)...	2.5	15(15)	9(7)	15(3)	9(0)	3(0)
Total of slight intermediate and normal (0.3).....	2.38	58(51)	34(29)	90(39)	48(1)	6(0)

TABLE 9 (continued)

(B) (Group B)

Grade of truncation of tested male parent	Average grade of truncation offspring	Truncate	Good intermediate	Intermediate	Slight intermediate	Normal
4 (truncate)...	2.5	26(22)	12(10)	31(15)	21(0)	0(0)
4 (truncate)...	2.5	6(6)	2(2)	11(9)	3(0)	0(0)
3 (good intermediate).....	2.4	29(23)	12(5)	32(17)	23(0)	3(0)
Total truncate and good intermediate (3.7)	2.45	61(51)	26(17)	74(41)	47(0)	3(0)
2(intermediate)	2.3	4(4)	1(1)	7(4)	4(0)	0(0)
2(intermediate)	2.4	24(24)	14(12)	28(15)	19(0)	3(0)
2(intermediate)	2.5	23(23)	13(11)	36(6)	14(1)	0(0)
Total (intermediate (2.)...	2.46	51(51)	28(24)	71(25)	37(1)	3(0)
1 (slight intermediate)	2.1	4(4)	2(2)	8(8)	8(3)	0(0)
1 (slight intermediate)	2.55	12(12)	8(7)	10(5)	10(0)	0(0)
0 (normal)....	2.9	18(18)	8(7)	13(3)	4(0)	0(0)
0 (normal)....	3.0	21(14)	3(2)	8(2)	7(1)	0(0)
Total slight and normal (0.5).	2.71	55(48)	21(18)	39(18)	29(4)	0(0)

between the grades of the parents and those of the offspring,—the truncate females (grade 4) yielded an average offspring grade of 2.3—, among their 147 gray red offspring; their normal sisters (grade 0) yielded a grade of 2.24, among their 107 gray red offspring.

On the basis of the various tests of sibs, then, the conclusion seemed practically inevitable that the variation in the truncate character, aside from that due to segregation of multiple heterozygous factors, is entirely "somatic," and not due to fluctuations in the genes.

b. *The "pure-line" experiment*

It was thought desirable to carry the tests of factor constancy further, however. For it might have been contended that the genetic variations in any one generation may be rather small, and as such inappreciable, but that they would have a chance to become evident if allowed to accumulate through several generations of selection. A scheme of selection was therefore undertaken, with this question in view. The selection had to be carried on in such a way, however, as to exclude the possibility of effects due to recombination and the sorting out of those combinations that tended in the direction of selection. This is usually the insuperable difficulty in the way of significant selection experiments in sexually reproducing, cross-breeding organisms; it is the flaw which invalidated the early conclusions on hooded rats, for example, and other similar experiments, in so far as their bearing on the question of factor variability is concerned. What is needed in such an experiment is the prevention of effective recombination (recombination which might have an effect on the progress of the selection experiment). This is the essential feature in the pure-line idea of Johannsen. But it is not really necessary for this purpose that the organisms be homozygous and self-fertilizing, as in Johannsen's definition of the pure line, or—what amounts to the same thing—that they be inbred until, according to the probability ratios worked out by JENNINGS, they have become homozygous. Neither is the only other alternative that they reproduce asexually, as in the work of JENNINGS and of a number of other investigators. There is a third method of pure-line work, which we wish to present here, that may be used in cross-breeding organisms like Drosophila, in which all the linkage groups are known. This method consists in exposing to view completely, in one parent at least, the process of recombination, by making use of "identifying factors," and then picking out, generation after generation, the same combination for selection.[5]

This process is really, in essential principles, the one which has been

[5] A useful, but not so efficient, modification of this is to carry on the plus and minus selection lines without reference to recombination, and then, at the end of the experiment, to "purify" the lines, or bring them to a common basis so far as modifying factors may be concerned, by means of crosses with a common stock containing the identifying factors. If there is any effect of selection still noticeable after the "purification," it must have been in the factor studied. This method is not so effective as that above, however, for the reason that it reduces the rigor of the selection with respect to possible variations in the factor under consideration, to have other variations, indistinguishable from these, occurring at the same time, which are due to recombinations of modifying factors.

already employed in the experiments just reported, where gray red flies having the same chromosome combinations were produced, and those with different grades of truncation were "selected" and bred, to find out whether they were genetically different. Here, however, the "selection" was carried on through only one generation. But in the cases where the gray red parents were males, it was possible to continue the selection further. For in these cases the gray red male offspring received (without crossing over) precisely the same paternal combination as their fathers had contained, and so they could be used for similar test crosses with black pink females from the stock bottle. As the latter crosses likewise produce offspring of the same type, these may again be used, and so, generation after generation, the selection might be continued indefinitely, by mating the gray red males to black pink females from stock. Such an experiment was accordingly undertaken with these flies.

It will be desirable, before examining the results, first to analyze the conditions of this selection experiment a little more closely. The original gray red males, that were to be tested, contained the chromosome combination $\frac{Y}{X}\ \frac{T_2'B}{b}\ \frac{T_3'P}{p}$. Of this combination, they transmit their Y, $T_2'B$, and $T_3'P$ chromosomes, without crossing over, to their gray red male offspring. The entire paternally derived equipment of the offspring is therefore a direct and unalloyed continuation of the paternal side of their father, at least so far as the first three chromosomes are concerned. The maternally derived chromosomes of the offspring, on the other hand—the X-, b-, and p-containing chromosomes—are not descended from the similar chromosomes of the father, but are received directly from the black pink stock, which was used as female parent. Thus these maternal chromosomes in the offspring have a chance of being slightly different from those which the gray red father contained, although they are really derived from the same black pink stock as those in the father. In the same way, when this cross is repeated in each generation the same paternal Y, $T_2'B$, and $T_3'P$ chromosomes are continuously handed down, but the X-, b-, and p-containing chromosomes of the father are always lost in the subsequent generation and a new lot of them is received from the black pink stock. It might at first sight appear as though this polygenetic origin of the maternal chromosomes might invalidate the pure-line experiment. However, all tests have indicated that the black pink stock is sensibly homozygous, so it is improbable that dif-

ferences in these maternal chromosomes would appreciably influence the characters of the offspring and thus interfere with the selection work. And much more important in this connection is the fact that since these chromosomes are discarded and renewed *at random* from the stock in each generation, any selection of factors in them (supposing factor differences do exist) can have no constant or cumulative effect on the progress of the selection. Any final and sustained differences in the effects of selection, on lines differently selected, would hence have to be due to changes which took place in that part of the germ plasm that has been handed down continuously in the selected lines—namely, the Y, T_2' B, T_3' P chromosomes.

The above conclusion, it should be noted, implies the assumption that there are no factor-differences for truncate, existing outside the three large chromosomes. For only the first three chromosomes are under watch in the present experiment, and while these are being handed down without recombination, as in a pure line, it must be remembered that any heterozygous factors in the fourth chromosome or elsewhere (i.e., in the cytoplasm) would have a chance of being sorted out and so influencing the final result of selection. The fourth chromosome could have been watched in the same way as the first three by crossing the flies to *bpb*$_e$ stock instead of to *bp*, and always choosing the gray red straight-winged male offspring as parents. If this had been done (as it was, in fact, for one generation) the transmission of the entire chromosomal genetic complex would then have been exposed to view and under control continuously. It was thought unnecessary to keep watch on the fourth chromosome, however, both because it is extremely small and also because, as has been previously mentioned, other tests had indicated there were no modifiers for truncate located there. It should, besides, be noted that such disregard of the fourth chromosome in the experiments could not invalidate the significance of a negative result. For, in case of a negative result (i.e., finding that there was no effect from selection), it would be proved, not only, (1) that all factors for truncate that had been included in the pure-line system (i.e., those located in the first three chromosomes, that had not been permitted to undergo recombination) had remained sensibly constant, but also, (2) that no factor-differences affecting truncate had *existed* outside this system,—in loci that had been allowed to undergo recombination. For if the latter factor-differences had existed, selection would have been effective by sorting them out. The same conclusion would apply both to factors in the fourth chromosome and to any supposititious factors (fac-

tor-differences) affecting truncate, that might be imagined to exist outside any of the four accepted chromosomes or linkage groups of Drosophila—that is, a negative result of the selection experiment would prove them non-existent.

The selection experiment was carried out in two lines—a "minus" or "low" line, and a "plus" or "high" line. The two lines were of course both derived from the same gray red ancestral male of composition $\frac{T_2'B}{b}\ \frac{T_3'P}{p}$ The count from this male (crossed by black pink female) is shown in table 10 (A), which gives as usual, only the gray red offspring of the cross. The average grade of truncation of these flies was 2.3+. One of the slight-intermediate males of this family was then chosen as the start of the low line, and one of the truncates for the high line, each being crossed to black pink females, according to the scheme explained above. The low line was then carried through twelve generations of selection, in each of which several gray red males showing the nearest approach to normal wings were chosen for parentage, and bred in separate bottles with black pink females from stock. In cases where there were conspicuous differences in truncation, between the families produced by these several males, that family giving the lowest grade of gray red male offspring was used to furnish the male parents of the next generation. The condition of this line after twelve generations of "minus selection" is shown in table 10 (B), which gives the distribution of truncation among the gray red offspring of the seven least-truncated males of this generation. It will be seen that the average grade of truncation for all these families together is 2.2+. The extremely slight difference from the original value, 2.3, is in the direction of selection, but in view of the much larger differences between families of the same line and generation, this figure must be considered as in extremely close agreement with the first finding.

The high line was carried on for thirteen generations. The method used was the same as in the other line, but of course the gray red males selected in this case were always of the highest grade of truncation available, instead of the lowest. It was found, as was to be expected, that in many cases no really high grade males were produced which could be used as parents in this line, owing to the sex-limitation of truncate and the fact that T_1' was not present; nevertheless the males chosen represented the highest grade of truncate that was being produced, and averaged about a grade higher than the males used as parents in the low

TABLE 10

Results of selection in the absence of recombination.
Only gray red offspring are recorded; total males and females not in parentheses, females in parentheses.

(A) Count from first cross of single gray red truncate male ($\frac{Y}{X}\ \frac{T_2'B}{b}\ \frac{T_3'P}{p}$) by black pink females, from which the two selection lines were derived.

Truncate	Good intermediate	Intermediate	Slight intermediate	Normal	Total	Average truncation
11(9)	5(5)	25(12)	10(0)	0(0)	51	2.33

(B) Counts from similar crosses of males of the "low line" after twelve generations of "minus selection."

Grade of tested male parent	Truncate	Good intermediate	Intermediate	Slight intermediate	Normal	Total	Average truncation
0 (normal)	28(28)	37(37)	30(28)	43(3)	27(0)	165	2.0
1 (slight intermediate)	29(29)	27(26)	32(15)	52(0)	16(0)	156	2.0
1 (slight intermediate)	23(23)	5(5)	20(6)	36(1)	0(0)	84	2.2
1 (slight intermediate)	26(25)	22(21)	32(10)	25(0)	0(0)	105	2.5
1 (slight intermediate)	12(12)	5(4)	11(0)	7(0)	0(0)	35	2.6
1 (slight intermediate)	17(16)	5(5)	4(1)	10(0)	0(0)	36	2.8
1 (slight intermediate)	10(9)	3(3)	8(3)	3(0)	0(0)	24	2.8
Total	145(142)	104(101)	137(63)	176(4)	43(0)	605	2.22

(C) Counts from similar crosses of males of the "high line" after thirteen generations of "plus selection."

Grade of tested male parents	Truncate	Good intermediate	Intermediate	Slight intermediate	Normal	Total	Average truncation
2 (intermediate)	9(9)	7(6)	30(18)	21(1)	1(0)	68	2.0
2 (intermediate)	8(7)	28(26)	49(24)	23(1)	0(0)	108	2.2
2 (intermediate)	19(15)	22(17)	45(19)	40(2)	0(0)	126	2.2
2 (intermediate)	14(14)	27(25)	25(13)	28(2)	0(0)	94	2.3
Total	50(45)	84(74)	149(74)	112(6)	1(0)	396	2.18

line. The condition of this line in the last generation, after thirteen generations of "plus selection," is shown in table 10 (C). Here there is close agreement between the grades of truncation in the different families, and the average of all turns out to be 2.2—. There is absolutely no genetic difference, then, between this line and the low line, in spite of a dozen generations of divergent selection. Neither of them, moreover,

departs significantly from the original value. This same agreement in genetic behavior was shown moreover not only in the final generations of the two lines, but throughout their entire history. We must accordingly draw the conclusion that the truncate character, usually so plastic and so very responsive to the touch of selection both in crosses and in inbred stock, and capable as it is of existing in all degrees of "potency," nevertheless remains absolutely unaltered by selection when all recombinations of its multiple heterozygous factors is rigidly prevented, in a form of pure-line experiment which enables the distribution of practically the entire chromosomal germ splasm to be placed under observation. In other words, the actual factors or genes for truncate, lying in the chromosomes, each had a definite value, and remained constant. The experiment at the same time demonstrates that no inheritance other than chromosomal is taking place, even in the case of a character genetically so complex and so elusive as truncate. That its inheritance was entirely confined to the chromosomal linkage groups—in this case, indeed, to the first three chromosomes—is proved by the fact that there were no hereditary modifications of the character possible, so long as only these chromosomal contributions remained the same.

Thirdly, the experiment disposes of the supposition, which certain geneticists might have advanced, that the factors for truncate may in heterozygous individuals be weakened, or undergo contamination by their normal allelomorphs. For here the truncate factors of both lines were kept heterozygous for twelve generations without their value or "potency" being affected.

THE CAUSE OF THE INCONSTANCY OF TRUNCATE STOCK

a. *Proof of the viability and non-sterility of flies homozygous for the intensifiers*

The question still remained unanswered, then, why had all the attempts to obtain genetically constant truncate stock ended in failure. Since the factors themselves were constant, and the stocks were inconstant, genetically, it followed that all the stocks must be, for some reason, heterozygous in one or more of the factors for truncate. This means that the homozygote, of one sex at least, must be either sterile, or nonviable, or else incapable of being formed at all. It was evident, however that this condition did not apply to all the factors for truncate, otherwise it would be possible, by out-crossing to normal, to obtain heterozygous truncates that were somatically and genetically of as high a grade of truncation as the best truncate flies of the stock bottle. Nearly

all the truncates resulting from out-crosses were, on the contrary, of conspicuously lower grade than their truncate parents. And it could be shown, more specifically, that the factor T_1' was not subject to this limitation. For, in the first place, T_1' could exist in fertile truncate males. This had been proved by the reciprocal crosses considered on p. 15 and 16. And as males which carry a given sex-linked factor in their single X chromosome are usually of a similar type, somatically, to the females homozygous for that factor, it was fairly certain that the females homozygous for T_1' were viable also. It was less certain that they were fertile, on account of the difference between the reproductive systems of male and female. The viability and fertility of females homozygous for T_1' were, however, independently proved by the fact that the daughters of the out-crossed truncate females were, on the average, just as frequently truncate, and of just as high a grade of truncate as the daughters out of the out-crossed truncate males (see page 16, and table 6). The males necessarily transmitted T_1' to all of their daughters; the females therefore must have done the same, and so must have been homozygous for T_1'.

It was also clear that not only T_1' but in addition either T_2' or T_3' (or, if they be compound, then one or more of the factors contained in them) could exist with some degree of fertility in homozygous condition. For the truncate or intermediate T_1' males derived from out-crosses of truncate females were usually of much lower grade than the truncate males of truncate stock. Since both alike might possess T_1' in their single X chromosome these differences in truncation were evidently due to the heterozygosis of the F_1 males in respect to some non-sex-linked factors (T_2' or T_3' or some component of them), for which the stock males were homozygous.

Crosses were then made to secure evidence, in the case of the two latter factors, as to whether or not they could exist in homozygous condition. For this purpose recourse was again had to crosses of single heterozygous gray red males $\frac{T_1'X}{Y}\ \frac{T_2'B}{b}\ \frac{T_3'P}{p}$ by stock black pink females. One of these crosses is shown in table 11(A); it is identical in nature with that of table 7, cross 3. From the offspring of this cross, flies were chosen for five parallel crosses, the results of which were to be compared.

The first three of these crosses were concerned with the question of homozygosis in respect to T_3'. The first cross was of a slight-inter-

TABLE II

Crosses to test the possibility of securing homozygous T_2' and T_3'.

(A) Preliminary back-cross of heterozygous truncate male to black pink stock

Nature of parents ♀	♂	sh T	Gray red T	I	S	N	Gray pink T	I	S	N	Black red T	I	S	N	Black pink T	I	S	N
$\frac{-}{-}\ \frac{b}{b}\ \frac{p}{p}$ Black pink long	$\frac{T_1'}{-}\ \frac{T_2'B}{b}\ \frac{T_3'P}{p}$ Gray red truncate		18 (0)	9 (4)	16 (16)	3 (3)			18 (3)	38 (17)				47 (21)				48 (23)

(B) Crosses of flies derived from (A)

		sh T	T	I	S	N	T	I	S	N	T	I	S	N	T	I	S	N
$\frac{T_1'}{-}\ \frac{b}{b}\ \frac{p}{p}$ Black pink long	$\frac{T_2'B}{b}\ \frac{T_3'P}{p}$ Gray red slight		10 (6)	18 (11)	11 (0)	3 (0)		8 (4)	14 (10)	19 (6)				40 (22)				41 (23)
$\frac{T_1'}{-}\ \frac{b}{b}\ \frac{T_3'P}{p}$ Black red long	$\frac{T_2'B}{b}\ \frac{p}{p}$ Gray pink long		5 (2)	20 (11)	14 (6)	5 (1)		8 (7)	20 (13)	23 (11)				34 (14)				43 (20)
$\frac{T_1'}{-}\ \frac{b}{b}\ \frac{T_3'P}{p}$ Black red long	$\frac{T_2'B}{b}\ \frac{T_3'P}{p}$ Gray red slight	11 (8)	7 (4)	4 (2)	7 (3)	1 (0)		3 (2)	3 (1)	5 (2)				24 (15)				7 (3)
$\frac{T_1'}{-}\ \frac{T_2'B}{b}\ \frac{p}{p}$ Gray pink slight	$\frac{b}{b}\ \frac{T_3'P}{p}$ Black red long			5 (3)	2 (1)	5 (2)			4 (4)	4 (3)	2 (1)	1 (1)	7 (5)					17 (8)
$\frac{T_1'}{-}\ \frac{T_2'B}{b}\ \frac{p}{p}$ Gray pink long	$\frac{T_2'B}{b}\ \frac{T_3'P}{p}$ Gray red slight	1 (1)	5 (3)	8 (4)	4 (2)	7 (3)		1 (0)	9 (4)	13 (7)	5 (2)			16 (8)		2 (1)		11 (3)

(The number of males is given in parenthesis under the number of flies of both sexes.)

mediate gray red male (composition $\frac{X}{Y}\ \frac{T_2'B}{b}\ \frac{T_3'P}{p}$) by one of its black pink sisters ($\frac{T_1'X}{X}\ \frac{b}{b}\ \frac{p}{p}$) derived from the same bottle. This

cross is of the same type as the parental one of table 11 (A) except that T_1' is distributed to half of the daughters and to half of the sons, instead of to all of the daughters and none of the sons. The count, given on line 1, table 11 (B), shows the results to be expected on the basis of the parental compositions, there being the usual four classes, gray reds, gray pinks, black reds, and black pinks, in remarkably equal numbers. Only the grays are truncated, and the gray reds especially, showing a count of 10 truncate: 18 intermediate: 11 slight-intermediate: 3 normal. ("Good intermediates" and "intermediates" are classed together in this and the following records.)

The second cross was like the first, except that the female was heterozygous for $T_3'P$, instead of the male. This cross therefore had the formula $\frac{X}{Y}\ \frac{T_2'B}{b}\ \frac{p}{p}$ (long-winged gray pink male) by $\frac{T'X}{X}\ \frac{b}{b}\ \frac{T_3'P}{p}$ (long black red female). It will be seen that the results, shown on line 2 of table 11 (B) are very similar in character to those of the first cross, except that there may be a slight decrease in the proportion of high-grade truncates among the gray reds, as compared with the first cross. This change, if not a chance deviation, is due to the crossing over between T_3' and P, which was possible in the female.

These first two crosses really served as controls for the third cross, which was concerned with the possibility of securing homozygous T_3'. The third cross was really a combination of the first and second, in that both the male and female parents were heterozygous for $T_3'P$, instead of the male alone, as in cross 1, or the female alone, as in cross 2. The parents, then, were a slight-intermediate gray red male which was a brother of the similar one used in the first cross and one of its black red sisters which was also a sister of the similar female that was used as a parent in the second cross. The formula for the cross is $\frac{X}{Y}\ \frac{T_2'B}{b}\ \frac{T_3'P}{p}$ by $\frac{T_1'X}{X}\ \frac{b}{b}\ \frac{T_3'P}{p}$. Here, since both parents are heterozygous for $T_3'P$, the reds should bear the ratio of 3:1 to the pinks, instead of 1:1, as before, and some of the reds,—one-third or fewer of them, depending on the exact amount of crossing over between T_3' and P,—will be homozygous for T_3', provided T_3' can exist in homozygous condition. This homozygosis in T_3', if it can exist, might then be reflected in a more strongly truncated condition of the flies of the gray red classes; if, on

the other hand, homozygous T_3' cannot exist, the gray red flies will not, on the average, be more markedly truncated than those of the previous cross. They would, if anything, average less truncate than in cross 1, because of the possibility here of crossing over between T_3' and P (red). Moreover, the ratio of reds to pink would be 2:1, or at least less than 3:1, if the reds homozygous for T_3' failed to appear. The count is shown on line 3 of table 11 (B). The ratio of reds to pinks is exactly 3:1 (54:18), as it should be if the homozygous T_3' live. Examining the gray reds, it is clear that there is among them a new class of individuals, with respect to truncate, that is never found in conspicuous numbers in out-crosses. This has been called "short-truncate"; it is of the same grade as the highly-truncated individuals of selected truncate stock. Almost exactly one-third of the gray red individuals belong to this grade. (It will be noted that the gray pinks, on the other hand, are no more truncated than before, there being no special environic conditions here favoring truncation.) It is therefore evident that the whole chromosome containing T_3' is capable of existing in homozygous condition, and that, provided T_1' and T_2' are present as heterozygous factors, then homozygosis in T_3' alone is necessary in order to produce an average grade of truncation equal to that of the best selected inbred stock. It was found, on attempting to breed the short-truncates produced in this experiment, that although the flies were not sterile, yet the infertility (of the females especially) was so great as to make it extremely difficult to perpetuate the line, just as in the case of the "best" inbred truncate stock. Crosses of the same type as that of table 11 (B), cross 3, have been repeated several times, and exactly similar results, including the production of a conspicuous class of short-truncates, have always been obtained.

The first, fourth and fifth crosses were concerned with homozygosis in respect to T_2'. The fourth cross was similar to the first except in that the female was heterozygous for T_2' instead of the male, so that here there could be crossing over between T_2' and B. The formula is $\frac{X}{Y}\ \frac{b}{b}\ \frac{T_3'P}{p}$ by $\frac{T_1'X}{X}\ \frac{T_2'B}{b}\ \frac{p}{p}$ (long black red male by slight-intermediate gray pink female). The parents were sibs of the parents used in the other crosses. This cross accomplishes the same purpose for T_2' that the second cross did for T_3'. The very small count, given on line 4 of table 11 (B) is nevertheless sufficient to show that the general character of the cross is like that of the first, except in that the grays here include fewer truncates, and that some of the blacks are truncated, owing to the

crossing over between T_2' and B in the female. The first and fourth cross served as controls for the fifth, in which opportunity was given for T_2' to become homozygous. In the fifth cross, both male and female (sibs of the previous parents) were heterozygous for $T_2'B$, instead of just the male, as in cross one, or just the female, as in cross four. This time, then, a slight-intermediate gray red male like that in the first cross was mated to a gray pink sister like that in the fourth cross ($\frac{X}{Y}\ \frac{T_2'B}{b}\ \frac{T_3'P}{p}$ by $\frac{T_1'X}{X}\ \frac{T_2'B}{b}\ \frac{p}{p}$). In this case, as both parents contain heterozygous $T_2'B$, the expected ratio of grays to blacks would be 3 : 1, instead of 1 : 1, as in the other crosses, and one-third or fewer of these grays (both reds and pinks) should be homozygous for T_2' and might therefore be more truncated than the corresponding classes of grays (red or pink) in the first cross. But if the grays homozygous for T_2' fail to appear, this 3 : 1 ratio should be reduced towards 2 : 1, and the grays in this cross should not be more truncated than those in the first cross. The count, on line 3, table 14, shows 48 grays to 34 blacks. This is nearer 2 : 1 than any other simple ratio possible here and certainly it could scarcely be taken for a 3 : 1 ratio. The gray reds, although including one short-truncate, such as appear in isolated instances in outcrosses to black, are not on the whole significantly more truncated than those in the first cross; the gray pinks, which are just as useful for the present purpose, are, if anything, less truncated than in the first cross, and do not include any flies that have a higher grade of truncation than that found among the gray pinks there. The two most truncated gray red males, moreover, were tested by crossing to black pink, and both proved to be heterozygous for T_2', giving both gray truncate and black long-winged offspring, and no black truncates. The evidence of these experiments hence indicates that T_2' cannot become homozygous. Yet it is not conclusive, for there is a possibility that homozygous T_2' is no more truncate in appearance than heterozygous T_2'. The deficiency in the number of grays, also, might be explained away, by supposing that special circumstances made all the grays here relatively less viable than the blacks. It should be mentioned, however, that crosses one and five were later repeated, with results similar to those of the present experiments: the ratio of grays to blacks in the repetition of experiment five was 61 : 32, and the truncate averaged of no better grade there than in the control cross corresponding to cross one (no short-truncates at all were produced this time).

b. *Experiments to test the possibility of securing homozygous* T_2'

Other crosses were then undertaken, which might determine more definitely whether the T_2' homozygote could occur. It would not have been feasible here to test flies for homozygosis in the ordinary simple manner,—that is, merely by finding the total proportion of truncates thrown, because the proportion of truncates is subject to modification in so many ways. A way was therefore found of "marking" the T_2' factors, by binding them to identifying factors (B and b), so that it could be determined in the tests whether a fly had only one T_2' factor, or two T_2' factors differently "marked", as will be explained presently. For this purpose, black truncate flies were secured that had arisen as crossovers in a cross similar to number four above (see line 4, table 11 (B). From these a stock of black truncate (inconstant for truncate, of course) was obtained. Black truncates (presumably heterozygotes containing $\frac{T_2' b}{b}$) were then mated to heterozygous gray truncates having the second chromosome composition $\frac{T_2' B}{b}$. Both reciprocal crosses were made. From these crosses gray truncate offspring,—usually the male offspring,—were chosen, and tested by crossing to black long-winged stock.

In the cross of $\frac{T_2' b}{b}$ males by $\frac{T_2' B}{b}$ females, the gray truncate male offspring that were tested may have received either a crossover gray chromosome,—B,—or a non-crossover gray chromosome,—$T_2' B$,—from their mother. In the former case, which would be less frequent, they must have received $T_2' b$ from their father, since they showed the truncate character; their composition, then, would be $\frac{B}{T_2' b}$. They would reveal this composition, in the test crosses with black long, by producing gray offspring ($\frac{B}{b}$) all long-winged, and black offspring ($\frac{T_2' b}{b}$) some of which were more or less truncated. (Enough intensifiers were present to insure the visibility of the truncate in most of the flies carrying T_2'.) Of the fifteen gray truncate males tested, four proved to have this composition, as they threw black truncates, but no gray truncates. Here there was no chance for homozygosis of T_2'. The other

eleven gray truncate males proved to have received $T_2'B$ from their mother, for they all gave rise to some gray truncates on crossing to black long. It was to be expected, if T_2' can be homozygous, that half of these eleven males would have received the $T_2'b$ chromosome from their father, the rest merely the b. The former, then, would throw black truncates, the latter only black longs, in addition to the gray truncates. As a matter of fact none of these flies which threw gray truncates threw black truncates, only longs appeared within the black class of offspring. The composition of all eleven must therefore have been $\frac{T_2'B}{b}$, although, if T_2' can occur in homozygous condition, half of them should have been $\frac{T_2'B}{T_2'b}$.

From the reciprocal cross, of $\frac{T_2'B}{b}$ male by $\frac{T_2'b}{b}$ female, 10 gray truncate male offspring were tested. In this cross, there will be no $\frac{B}{T_2'b}$ class, for all the gray offspring must receive $T_2'B$ from their father, since there is no crossing over between T_2' and B in the male. Correspondingly, it was found that all the (gray) males tested threw gray truncates. Half of these males, as before, may have received b from their other parent, and the rest $T_2'b$. But, as before, it was found that all the black offspring were longs; here again, then, all the flies had been heterozygous, $\frac{T_2'B}{b}$, although there had been a chance for the half of them to have been homozygous, $\frac{T_2'B}{T_2'b}$. Five of the gray truncate female offspring of this cross were also tested, by crossing with black long males; similar considerations apply to them, except that a heterozygous $\frac{T_2'B}{b}$ female would throw some black truncate offspring by crossing over; it would not throw nearly as many black truncates as gray truncates, however, and so it could be distinguished from the homozygous $\frac{T_2'B}{T_2'b}$ female. It was found that all five females were of the heterozygous type, $\frac{T_2'B}{b}$, just as in the case of the males.

Another cross, similar in principle, involved the dominant identifying factor star, in chromosome II, as well as black. A black pink truncate fly (obtained by crossing and subsequent selection) was crossed to star pink. A son showing the characters star, truncate and pink,—composition $\frac{T_2' b}{S' B}\ \frac{p}{p}$, —was then crossed by a short-truncate female from the inbred truncate stock. Forty-six male offspring which did not have star eyes, and which had therefore received $T_2' b$ instead of $S' B$ from their father, were then tested by back-crossing them separately to non-truncate black pink females. The composition of these males must have been $\frac{Y}{T_1' X}\ \frac{T_2'\quad b}{T_2'(?)B}\ \frac{p}{T_3' P}$; the question was whether any of them had received $T_2' B$ from their truncate mother, in addition to the $T_2' b$ that they must have gotten from their father. If T_2' can be homozygous, at least half should have received $T_2' B$, and should therefore give rise to gray truncate offspring as well as black truncates. All forty-six, however, failed to produce any gray truncates, although they all threw black truncates in very high percentage, particularly among the red female offspring, which contained T_3' and T_1'. The original short-truncate female from selected stock must therefore have been heterozygous for T_2', and when her eggs were fertilized by the T_2' b-bearing sperm, only those which had failed to receive the maternal T_2' developed. Adding these 46 to the previous 26 tests, there is a total of 72 flies, at least one-half of which should have been homozygous for T_2', had this been possible, but all of which were found to be heterozygous.

We may therefore conclude with certainty that homozygous T_2' cannot exist. This explains not only why there was such difficulty met with in the attempt to get pure truncate stock, but also why truncate, even when it acted as a dominant in crosses with black, never gave more than half of the F_1 showing truncate (see tables 6 and 7). Twin hybrids, truncate and long, were produced, as in out-crosses of Oenothera, Matthiola, and beaded Drosophila (see page 41), half of them in the present case receiving the T_2' chromosome and half the t_2'. Tests of such F_1 longs, moreover, showed many of them to be genetically long, and not merely T_2'-carrying flies that lacked intensifiers or happened to vary somatically in the direction of normal (although of course, some of the longs were of this type, in cases where more than half of the F_1 appeared long). Table 12, lines 1 and 2, shows the result of such a test of an F_1 long male which had arisen from an out-cross of a short-truncate female

of inbred stock by a black pink male. When this long was back-crossed to black pink, many of the gray red offspring should have appeared truncate, had the F_1 carried T_2', but, as the table shows, not one truncate appeared. One of the slight intermediate F_1 males, on the other hand, which was a brother of the long, gave the count on the second line, when tested similarly. Thus, even the short-truncate of inbred stock is heterozygous for T_2', and gives the results expected of a heterozygote, on out-crossing. It may be noted that the tests of these normal-appearing F_1 males are in striking contrast to the results obtained by crossing normal-appearing males from a back-cross like that in table 7, cross 3, where

TABLE 12

Tests of F_1 males from a cross of short truncate female by black pink male. F_1 males tested by back-crossing to black pink.

Character of F_1 male tested	Gray red				Gray pink				Black red				Black pink			
	T	I	S	N	T	I	S	N	T	I	S	N	T	I	S	N
Normal	0	0	0	21	0	0	0	25	0	0	0	23	0	0	0	26
Slight intermediate	0	3	3	2	0	0	2	14	0	0	0	17	0	0	0	11

the identifying factors show that the flies, though normal, must really carry the factors for truncate.

c. *The reason for the non-appearance of homozygous T_2'*

The failure of T_2' to appear in homozygous combination must be due either to the fact that it acts, when homozygous, as a lethal factor, or that there is a selective fertilization of T_2' eggs by t_2' sperm exclusively. In the latter case, when heterozygous $\frac{T_2' B}{b}$ flies were crossed with each other, a ratio of 3 grays ($\frac{T_2' B}{b}$) : 1 black ($\frac{b}{b}$) should be produced, since all $T_2' B$ eggs would be fertilized by b sperm, and half the b eggs would be fertilized by $T_2' B$ sperm. In the case of a lethal factor, on the other hand, only 2 grays: 1 black should appear, for one-third of the grays would consist of the homozygous $\frac{T_2' B}{T_2' B}$ combination, that would die. We have seen (page 35) that the 2:1 ratio is actually obtained; thus T_2' is a lethal, like yellow in mice, and like most dominant mutant factors, in Drosophila at any rate.

d. *The cause of the infrequency of normals in the selected stock*

This "yellow-mouse" hypothesis had of course often been considered, as a possible explanation of why stocks of truncate were always genetically inconstant, but it had not previously been thought probable, because the highly selected, inbred truncate threw only 10 percent of normals (and many of these, even, carried T_2') whereas yellow mice throw as much as 33⅓ percent of normals (or non-yellows). A cross of $\frac{T_2'}{t_2'}$ by $\frac{T_2'}{t_2'}$ likewise should give 2 $\frac{T_2'}{t_2'}$ (more or less truncated) : 1 $\frac{t_2'}{t_2'}$ (long), provided the $\frac{T_2'}{T_2'}$ die. Either, then, in the case of the selected truncate, most of the $\frac{t_2'}{t_2'}$ which are formed die also, owing to the existence of another lethal or semi-lethal factor in the chromosome with t_2', or, if they live, many of them must appear truncate, like the $\frac{T_2'}{t_2'}$ flies, owing to some other chief factor for truncate that they may contain. This other factor would have to lie in the second chromosome, for we have seen that flies containing other chromosomes of truncate stock, but not having the second so derived (the black red flies of table 7, cross 3), even when inbred, cannot give rise to truncates (table 7, cross 5). It would, moreover, have to be a recessive, for we have seen that heterozygous flies which fail to receive T_2' are never truncate, even if one of their second chromosomes is derived from truncate stock (p. 39).

A test was therefore undertaken, to determine whether there existed, in the t_2'-bearing second chromosome of the selected stock, a recessive factor for truncate, which, when homozygous, was able to cause the truncate character to appear, even in the absence of T_2' itself. This test was performed by inbreeding the gray red flies ($\frac{T_1'\,X \text{ or } Y}{X}\ \frac{t_2'\,B}{b}\ \frac{T_3'\,P}{p}$) obtained in the last tests for homozygous T_2', reported on p. 38. All these flies had received, it will be remembered, one gray-bearing chromosome derived from a short-truncate female of the selected stock; this chromosome had been found not to contain T_2'. If, however, it contained a recessive "chief" factor for truncate, in any of the families (the chromosome was not necessarily the same in the different families, on account of the possibility of crossing over in the stock truncate fly from

which it was derived), then, when the gray flies of such a family were crossed together, some flies homozygous for the recessive truncate would be produced, and many of these should show the truncate character (since the intensifiers also are present). Nineteen of these tests were made in all, the gray red offspring of each of nineteen of the males that had been tested for homozygosis being inbred to each other in mass culture, family by family; thus, if the original truncate female had been heterozygous, so that only certain families contained the factor, it would have a chance to become homozygous in these lines. None of the 19 families produced any truncates, however, and the hypothesis of a recessive "chief" factor to explain the low proportion of normals in the selected truncate stock therefore had to be discarded.

The only remaining possibility was that the truncate stock contained a lethal (or semi-lethal) factor in the second chromosome that carried t_2', for since it was shown that a $\frac{t_2'}{t_2'}$ combination in the selected stock would not give rise to truncate-winged flies, the excess of truncates (or deficiency of longs) proved that there was a real deficiency, i.e. a dying off, of the $\frac{t_2'}{t_2'}$ flies that were formed at fertilization.

A similar condition of affairs had been found by MULLER (1917) to exist in the case of the beaded-winged stock of Drosophila, and was called by him a state of "balanced lethal factors." In such a case, there are present, in homologous chromosomes, two different lethal factors, each of which kills off an opposite type of homozygote, and which, together, thus cause a heterozygous stock apparently to breed true, or nearly true. The balanced lethals in the case of the truncate stock are T_2', which kills the homozygous truncates, and the recessive lethal factor in the opposite chromosome, which kills the homozygous $\frac{t_2'}{t_2'}$ flies.

As shown in the report on the beaded case, the occurrence of such recessive lethals by mutation must be comparatively frequent. Bearing this in mind, the existence of this factor in the truncate race is easily explained, as follows: A stock having a character such as truncate, which is itself due to a lethal (T_2'), must continually throw normals. When such a character is selected, therefore, an endeavor will be made to select a line throwing as few normals as possible. In the course of time, various lethal factors will arise, in different loci, most of which will be eliminated by natural selection, but when a lethal arises in the

normal homologue of the chromosome containing the factor that is being selected for (T_2'), it will cause the killing off of a large number of the undesired normals; consequently the line in which this factor has arisen will be perpetuated in the selection. The nearer the locus of this factor lies to the locus of the desired factor (T_2'), the fewer will be the normals that can escape, by crossing over, from the lethal action, and so the better will be the chance of the perpetuation of this race in the experiment. After a long course of selection, then, it is to be expected that the stock would show the presence of a lethal factor nearly opposite, in location, to the lethal factor that is being selected for, and a condition of "balanced lethals" would have arisen. For the further consequences of this condition the reader must be referred to the account of the beaded case.

e. *The source of the normals which did occur in the selected stock*

In the case of the truncate stock, it is likely that there was a certain amount of crossing over between T_2' and the recessive lethal. This would partly explain the presence of the 10 percent of normals in the selected stock, for such normals could arise from a $\frac{T_2' L}{t_2' l}$ female, from a crossover gamete $t_2' L$ (l here indicates the recessive lethal; L its normal allelomorph). Such normals would never give truncates if crossed together, as they do not contain T_2'.

Table 2 showed, however, that some of the normals thrown by the stock could give rise to truncates. Some of these normals, then, may have been merely somatic variants towards normal, which nevertheless had the full complement of truncate factors. In that case it would have to be assumed that in these experiments they had been mated to the crossover normals, which lacked T_2' and which would be indistinguishable from them somatically, for these matings gave fewer truncates than ordinary matings of somatic truncates. But it seems unlikely that this special sort of mating should be made frequently, and so it is more probable that most of these normals, which were able to throw some truncates, were T_2'-bearing flies that lacked one or more of the intensifiers. It has been explained that females homozygous for the intensifier, T_3', are extremely infertile, and since in the selected stock even those eggs which are laid are killed off in large numbers by the lethals of chromosome II, it will be readily understood that truncate stock homozygous for T_3' could not be long maintained. It was, in fact, found continually

in the original selection experiment that the very high lines died out through infertility, and that the stock had to be replenished from mass cultures of somewhat lower grade. The stock consequently was probably impure for T_3' as well as T_2', even though many of the individuals in it must have been homozygous for the former factor.

THE LOCUS OF TRUNCATE

The above experiments were sufficient to demonstrate the nature of the mechanism of inheritance of the truncate character. They showed that it conformed regularly to the principles of chromosome heredity, factor constancy, etc., and disclosed the causes of the inconstancy of the inbred stock, and of the indefiniteness and variability of the ratios thrown in crosses.

Further determination of the precise number of modifying factors for truncate present in each chromosome, and the location and exact effect of each, in the various possible combinations, would have required, for the most part, elaborate tests—much more elaborate than the preceding ones—and yet these details would have been of little general significance, once it had been found out what was really "the trouble" in the truncate case. Some more exact tests were at first undertaken, with these objects in view, but many identifying factors had to be used for this purpose, in order to follow the crossing over, and it was found that some of these factors also influenced the truncate character. This circumstance, taken together with the fact that the classes containing the intensifiers far overlapped those lacking them, on account of the natural somatic variation of truncate, made a direct estimation of the percentage of crossovers impossible. An example of the confusion caused in these ways is seen in table 13, which represents a count of 196 offspring of a back-cross designed to find the locus of T_3'. A female heterozygous for truncate and for three third-chromosome identifying factors—pink eye (p), kidney eye (k), and sooty body color (e^s), was back-crossed to a pure $p\ k\ e^s$ male. It will be seen that the truncate shows rather close linkage to P, and yet the crossover classes all show a fair amount of truncate, which is distributed among them nearly proportionally. Thus the truncate intensifier shows no closer linkage to P than to either of the other factors, and its locus is indeterminable. Such a result can be explained only by supposing that several modifiers of truncate are here concerned.

Difficulties of the above sort would not have been insuperable, as the flies from such a count could each be tested out, separately, but a much

TABLE 13

Back-cross of truncate to pink kidney sooty.

$$\left(\frac{T_1'\ T_2'\ T_3'}{pke^s}\ \text{♀} \times \frac{p\ k\ e^s}{p\ k\ e^s}\ \text{♂}\right)$$

	P K E				*p k e*^{s}				*P k e*^{s}				*p K E*				*P K e*^{s}				*p k E*			
	T	I	S	N	T	I	S	N	T	I	S	N	T	I	S	N	T	I	S	N	T	I	S	N
♀	3	1	0	34	1	4	2	36	1	1	0	1	0	0	0	8	0	0	0	6	0	0	0	4
♂	3	4	1	30	0	4	0	32	0	0	0	4	0	1	0	2	0	1	1	7	1	1	0	2

more elaborate system of experiments would have been necessary for this purpose than the value of such information justified. There would, moreover, have been no such thing as a "complete account" of the truncate case, unless all known and unknown mutant factors of Drosophila had been tested, in all possible combinations, with respect to their effect on truncation. It was therefore decided to confine the "intrachromosome" investigation simply to a determination of the locus of T_2', the chief factor for truncate.

Certain data regarding this may be obtained from the counts of table 7(B), crosses 4, 6, and 9, where females heterozygous for truncate and black ($\frac{T_2'\ B}{b}$) were back-crossed to black long males. Here the long-winged offspring must be disregarded, as they are genetically heterogeneous, including many flies that carry T_2 besides the really genetic longs. The percentage of crossing over may, however, be obtained from the count of those that show any degree of truncation. Among the latter, the grays are of course non-crossovers, the blacks crossovers. There are 173 flies in these three crosses showing some grade of truncation, and among them 41 are black (crossovers), the rest gray. This would make the crossing over between truncate and black $\frac{41}{173}$, or 24 percent. In various crosses of the same type 375 truncates have been counted in all, including 100 blacks; this makes the percentage of crossing over nearly 27.

Truncate was also crossed to a stock of black purple curved (all in chromosome II), and F_1 truncate females were back-crossed to $b\ p_r\ c$ males. The count of 530 individuals is shown in table 14; there is the usual percentage of crossing over between b, p_r and c (b-p = 6 percent, p_r-c = 17 percent). To find the linkage with truncate, only the 119 off-

spring showing truncation can be used, just as before. Among these there are 22 percent of blacks, indicating a "distance" of 22 units in this cross. This, averaged with the previous results, would place T_2' at 25.5 units from black. Double crossing over is so free in this cross, however, that it is not possible to tell with certainty whether T_2' is on the same side of black as curved is (i.e., to the "right") or on the opposite side (the "left"). Yet the fact that in crossing over between black and purple the truncate remained with the black, indicated that T_2' was probably on the opposite side of black from purple,—that is, to the "left".

A test was accordingly made to determine on which side of black T_2' lay. This was done by means of a back-cross involving truncate, black,

TABLE 14

Back-cross of truncate to black purple curved.

$$\left(\frac{T_1'\ T_2'}{b\ pr\ c}\ ♀ \times \frac{b\ pr\ c}{b\ pr\ c}\ ♂\right)$$

	BPr C				*bpr c*				*Bpr c*				*bPr C*			
	T	I	S	N	T	I	S	N	T	I	S	N	T	I	S	N
♀	26	13	6	56	7	5	0	69	0	1	0	3	0	0	0	10
♂	20	12	11	83	5	1	2	99	0	0	0	4	0	0	0	8

	BPr c				*bpr C*				*Bpr C*				*bPr c*			
	T	I	S	N	T	I	S	N	T	I	S	N	T	I	S	N
♀	0	0	0	10	3	2	0	25	1	1	0	1	0	0	0	2
♂	4	0	0	10	0	0	1	30	0	0	0	1	0	0	0	0

and star. Star (S') is a dominant factor lying 46 units to the left of black, at locus 0. In the first place, a star truncate stock was secured by crossing truncate with star and selecting the crossovers in the second generation, and in the second place this star truncate was crossed to black, and one of the heterozygous female offspring, which appeared star truncate, was back-crossed to a black male (triple recessive). The count, given in table 15, shows clearly that T_2' is to the left of black, since twenty-three of the twenty-four flies known to be crossovers between truncate and black (i.e., those showing both truncate and black) are also crossovers between star and black (star, truncate, black). The exact percentages are not very significant in a count of this size (total count 130) but the order of the factors is determinable with certainty.

When all these tests of truncate with black are averaged together, a

TABLE 15

Back-cross of star, truncate and black.

$$\left(\frac{T_1'}{\quad}\ \frac{S'\,T_2'}{b}\ \frac{T_3'}{\quad}\ \varphi \times \frac{b}{b}\ \frac{T_3'}{p}\ \delta\right)$$

Non-crossovers		Crossover 1		Crossover 2		Double crossover 1, 2	
Star truncate	Black	Star black	Truncate	Star truncate black	Normal	Star	Truncate black
41	36	6	6	23	14	3	1

percentage of crossing over of about 27 is obtained. The distance between truncate and star, as given in the last test, is 12.3. As the percentage of separations between star and black is known to be about 39, the results of the tests of truncate with star and with black agree fairly well, and it may be concluded that truncate is between star and black, at a locus of about "12"—i.e. 12 units to the right of star, which is taken as being at 0.

ON THE MODE OF ORIGIN OF THE TRUNCATE COMPLEX

On the basis of the information that has been gained concerning the genetic basis of the truncate complex, its mode of origin may in a general way be reconstructed. It is probable that either the factor T_2' or one of the intensifiers had arisen by mutation and persisted unnoticed in the ancestral beaded line for some time before the first truncate fly was discovered. For the original truncate, it will be remembered, gave rise to a moderate proportion of truncate F_1, and to a much higher proportion of truncates in some of the F_2 families. Now, the moderate proportion of truncates in F_1 would not have been possible in the first place, had not an intensifier of truncate been present,—it is only necessary to recall here the extremely small proportion of truncates produced in F_1, in the absence of black, even when the intensifiers are present in heterozygous condition (see table 3(A), representing crosses of selected stock to wild type). Secondly, had not an intensifier been present, it would not have been possible to raise so materially the proportion of truncates in F_2, since only intensifiers, and not T_2' itself, could have been made homozygous by this process.

One possibility, then, is that T_2' had existed for some time in the stock, and suddenly burst into visibility when T_1' or T_3' arose by mutation. Before this occurrence, it would have been detectable practically only as

a lethal factor, by the disturbances it produced in ratios, or, if elaborate tests were made, as a factor giving the fly a very slight "instability" of wing shape, or "tendency" to have shortened wings, which manifested itself once perhaps in 100 cases ordinarily, but much more often in the presence of the factor for black (thus recalling the alleged mode of inheritance of certain nervous defects in man). On the other hand, it is more probable that one of the intensifiers had existed first, for these would have been better able to persist in the stock than a lethal factor. The intensifier would not have been detectable at all at this time, unless, in the case of T_3', tests of fertility had been made. But, whichever of the two suppositions is true, it is evident that one of the mutant factors had arisen and persisted unobserved until the second mutation took place.

This situation raises the question as to how many such mutations—producing no visible result on the organism, but changing only the composition of the chromosomes, and, through them, the ultimate potentialities of the race—may be continually occurring unseen. Certainly the occurrence of such mutations is by no means rare, judging by the comparative frequency with which modifying factors are found, in races which happen to have the "chief" factor for that character already present. Thus, when selection is made for bar eye, bar modifiers are found; on selection for beaded wings, beaded modifiers; on selection for dichaete bristles, dichaete modifiers; all of which modifiers have little or no effect on the fly lacking the chief factor for bar, beaded, or dichaete, respectively. Yet such mutations must be occurring with similar frequency (though not selected for and artificially perpetuated) in stock lacking the specific chief factor; so, in wild stock, modifiers must be arising for bar as in the bar stock, and at the same time for beaded as in the beaded stock, for truncate as in the truncate stock, and similarly for innumerable other potential characters not actually realized in the stock at that time. Hence it may well be that there are more mutations for which no means of influencing the somatic characters exists, than visible "character-mutations," and undoubtedly there are many on the edge of visibility. Many of these "invisible" or "barely visible" mutant factors should persist in particular lines, merely as a matter of chance, and they would affect not only chromosome composition and behavior, but also the possibilities of future character-evolution for that line. Such, at least, was the case with the truncate line which, by means of the prior possession of an "invisible" factor for truncate, was enabled to produce the truncate "evolution series." Here is a situation which may give some satisfaction to believers in orthogenesis.

The truncate which had thus appeared probably contained, as yet, only one intensifier, and no balancing lethal, for it could not be increased by selection for a considerable time. When, however, each of the other contributing factors finally appeared in the course of the breeding, the line containing it was purposely selected, since it threw more and better truncate. It has been previously explained that there is nothing remarkable about the fact that a balancing lethal did arise. The fact that it was also possible to find one or more additional intensifiers may however call for some comment. This responsiveness of truncate to selection,—the appearance of intensifiers when sought for,—may be shown to be an inherent characteristic of the truncate race, dependent on a particularly high modifiability of the embryological process whereby the truncate character is produced. The degree of modifiability of the developmental reaction that causes any given character may be estimated by at least two criteria: (1) by the somatic variability of the character among individuals of the same genotype (a test of responsiveness to environmental influences); (2) by the proportion of previously known factors for other characters which are able to influence the character under consideration (a test of responsiveness to genetic influences). To both of these tests truncate gives a positive result, for we have seen, (1) that, when flies carry T_2', the wing shape is unusually variable even among individuals of identical factor composition, and (2) that, among these flies with T_2', a number of factors, like that for black body color, star eyes, bar eyes, sex, etc., which ordinarily produce no noticeable effect upon the wing, now influence its shape decidedly. Hence the developmental reactions that determine the wing shape must be particularly unstable, or susceptible to modification, in the race containing T_2'. From this conclusion in turn it follows that relatively many modifiers of wing shape should be found to arise by mutation in the truncate race,—not because mutation itself happens oftener in this race, but because, of the mutations that do occur, more would be able to affect the wing character in this race than in non-truncate races, where wing shape was not so easily influenced. A race of this kind, accordingly, would be able to undergo an evolution or cumulative change, under the directive action of selection, which would be next to impossible in another race, that lacked the factor causing this embryological modifiability. Here again, then, we have an example of invisible peculiarities of the germ plasm of certain lines, influencing the possibilities of character-variation and evolution.

The question might here arise—if characters may differ so considerably in regard to the modifiability of their development reactions, is it

not conceivable that certain characters may be so highly modifiable,—influenced by such a large proportion of the total number of mutations that occur,—and that this *total* number of mutations is at the same time so large, that, even though each separate single factor of the germ plasm is highly constant, nevertheless these characters would, in effect, really be undergoing a continual genetic fluctuation. This contingency cannot be categorically denied, as an academic possibility for exceptional cases, but we have seen that in the case of truncate, although it is a particularly unstable character, yet the number of mutations which affected it were not nearly so frequent as would be required for such an effect,—the pure line remaining constant through twelve generations. Of course, if selection had continued indefinitely, some distinct genetic changes affecting the character would have been bound to occur, but not of the order of frequency of "continuous variation" (fluctuation). The reason that the inbred stock, on the other hand, was so much more plastic genetically is simply because there was in it an opportunity for the formation of all sorts of genetic types by mere recombination of the various mutant factors that had already arisen at widely separated occasions. The inheritance of truncate therefore provides no evidence even for this modified type of genetic fluctuability which might appear as the resultant of innumerable mutations in many different factors, each of which itself was highly constant. Moreover, no other cases affording evidence of such a situation have been reported thus far.

We may now sum up the causes why the truncate character, when it was analyzed, was found to depend on so many factors. (1) Two successive mutations were practically necessary, in the first place, before truncate became visible at all. (2) Since truncate depended on a lethal, and was, in addition, inconstant in somatic expression, it was subjected to selection, which perpetuated any new factors (balancing lethals and intensifiers) that still further differentiated truncate from normal. (3) Since the truncate character depended on an unstable developmental reaction, the initial appearance of such intensifiers was made more likely.

The history of the truncate stock, after the appearance of these contributing factors, consisted merely of the attempt to get the line as pure as possible for T_3', and to keep the line in the balanced lethal condition by eliminating crossovers.

THE ORIGINATION OF TRUNCATE IN OTHER STOCKS

It is not very uncommon to find truncate appearing sporadically, like a mutant, in other stocks. Thus, in DUNCAN'S (1915) search for mu-

tations, truncate was the second most common in occurrence, beaded arising most frequently. Now, if the above account of the origin of the truncate complex has been correct, such truncate should not be provided with the accessory factors present in the selected stock, or at least not with the same ones. That this is true is indicated by the fact that often, when these truncates are bred, they fail to reappear, except in trifling proportions, in subsequent generations. The latter flies, then, if they owe their truncation to the same factor as the T_2' of truncate stock, evidently lack all the intensifiers, just as we should expect to be usual in stock not subjected to selection. In this connection, it may be noted that since the intensifiers are missing it is by no means certain that all these appearances of truncate represent separate mutations, as the factor for truncate may have been present for a long time, under such circumstances, without manifesting itself. Hence the actual mutations may really be rather rare.

Truncate has also appeared several times in stock containing factors for other characters, that at the same time intensify truncate, and these cases may with more certainty be regarded as real mutations. The composition of such truncates is accordingly more worthy of examination. One such truncate was found by MORGAN in a stock with "cut" wings; special crosses proved that it contained T_2', identical with the T_2' of truncate stock. Cut, itself, intensified the truncate, but there was no evidence of the regular intensifiers in addition. Another truncate was found by the writers in a cross between black and a stock that had been continually tested by crossing to bar. The mutant, a "long-truncate" female, was crossed to black pink, and gave offspring half of which were intermediate and the rest normal-winged. Two of the heterozygous intermediate F_1 males were then back-crossed separately to black pink. As in the similar crosses of ordinary truncate, truncate appeared only among the gray offspring (F_2), but it appeared about as often among the gray pinks as among the gray reds, giving in both classes the percentage and grade ordinarily present among the gray pinks. Thus it was evident that the intensifier T_3' was not present. The T_2' was evidently the same as that in ordinary truncate stock, for when the gray pinks ($\frac{T_2' B}{b} \frac{p}{p}$) were inbred they produced 81 grays and 51 blacks, instead of a ratio of 3:1, thus showing that this T_2' also was lethal. There was, moreover, a fairly large percentage of crossing over between black and truncate.

We may accordingly conclude that the origin, by mutation, of the intensifiers present in truncate stock, was quite independent of the origin of T_2' itself by mutation.

APPLICATIONS OF THE METHODS

It is believed by the authors that the general method of attack developed in the truncate case,—whereby, by the use of "identifying" genes, a refractory character may be taken apart, put together, or held in a desired combination—will become of more widespread applicability as the linkage groups of the organisms commonly used for genetic study become better known. Since a large part of the work on truncate was carried out nearly seven years ago this general method has already had a chance to be tried out rather extensively in Drosophila work, and it has been used with success by students of this organism in a considerable number of problems, some of which have been reported in the literature.

In the case of other organisms, however, it will be necessary to know and have the use of identifying factors in the various linkage groups concerned, before the present method can be used at all. Moreover, certain modifications are necessary in the case of organisms that have crossing over in both sexes, and also in case the character to be studied is recessive. In either of these circumstances, it will be desirable to have two identifying factors in each chromosome involved, one on each side of the factor under investigation,—in order to make sure that the factor has not crossed over from its identifier.

Remote as the possibilities of such work may seem to be in the case of such animals as mammals, it is nevertheless difficult to conceive how the genetic bases of the more elusive and complicated characters in them can be determined adequately by any other means. Even in the case of man, an attempt in such a direction would be justified, for here the most important characters,—such as the psychological ones,—are perhaps more plastic, obscure, and complicated in genetic basis, than any others in the entire animal kingdom, and it would seem next to impossible ever to give any real Mendelian analysis of most of them without studying them by the method of linked identifying factors. In place of controlled crosses, and large families, however, reliance would here have to be placed on finding, by means of wide examination of data, a sufficient number of crosses of similar type, and then seeking in them the requisite identifying factors.

It would accordingly be desirable, in the case of man, to make an ex-

tensive and thorough-going search for as many factors as possible that could be used in this way, as identifiers. They should, preferably, involve character differences that are (1) *of common occurrence,* that are (2) *identifiable with certainty,* and that are (3) *heritable in a simple Mendelian fashion.* It seems reasonable to suppose that in a species so heterozygous there must really be innumerable such factors present, if only an examination of the inheritance of *small, definite physical traits** were made on a large scale. As the study of such factors should naturally be accompanied by an examination of their inheritance with relation to each other, a knowledge of their grouping would at the same time gradually become available. All this would of course require very detailed and intensive work (rather, perhaps, than a superficial study of numerous individuals), and as yet little work has been done that is of this character. For, hitherto, the study of factors which are inconspicuous, or unimportant in actual life, has been largely avoided, in order to make an immediate and direct attack on the more important, more difficult characters. Now, as any two parents of a human family would probably differ nearly always in a very large number of factors, it would not be at all surprising if it were found that there were usually differences in one or more identifying factors in the case of any given one of the twenty-four (±) pairs of chromosomes. Thus, if a far-reaching investigation of definite physical traits were carried out first, then, when investigation of the more difficult characters was later undertaken, the requisite identifying factors, suitably arranged in the appropriate chromosomes, would probably be found, in many crosses, ready provided for the study of the more complicated and important trait.

The implication is not intended that no results of importance in human genetics can be attained by simpler methods; it would of course be desirable to carry on such studies at the same time as the more rigorous ones, but the inconclusiveness of the pedigrees of most important human characters, when studied directly, and their resemblance to the early pedigrees in the truncate case, indicates that nothing like an adequate understanding of the intricacies of inheritance in man can be reached without some such far-reaching and difficult plan as that just outlined. Meanwhile, too, partial knowledge of "identifiers" would be of partial help, in the larger problems.

However the situation may be for human genetics, it does seem clear that in the more tractable organisms, such as the domesticated and laboratory races of animals and plants, character analysis by means of linkage studies with identifying factors will come into more general use,

* Or chemical, such as the blood agglutinins.

both for the investigation of general problems and for the dissection of particular characters.

RESUMÉ

The general hereditary behavior of truncate

1. Truncate is a variable character, appearing in all grades between short-truncate and normal wing. Some normals are almost always thrown, even by the "best" truncates, and it was found impossible, through four years of selection, to secure a permanent stock that threw no normals, although the proportion of the normals was reduced to about 10 percent, and the average grade of the truncates which did appear was increased markedly at the same time.

2. This variation is not only somatic, but also genetic, for in the selected truncate race the high-grade truncates that appear throw relatively more and "better" truncates than do the intermediates, and the latter in turn greatly surpass the normals in this respect. It is thus possible, in the final stock, to modify the average grade and percentage of truncates back and forth by selection, within the limits above stated.

3. When truncate is crossed to the wild-type fly it behaves as an incomplete recessive, the great majority of F_1 being normal, but a small percentage of flies that show some truncation usually appearing also. The ratios in F_2 vary greatly with individual pairs of F_1 flies, ranging from nearly four normals to one with truncation through various values down to over 100 normals to one truncate, while some of the F_2 families contain no truncates at all. The percentage of truncates in F_2 is higher if the P_1 truncate was a female, but in either case the truncates in F_2 consist of both males and females in comparable numbers. The extracted F_2 truncates throw about two normals to one truncate but can often be improved by selection until they reach the limit shown by the selected stock.

4. It was found that on crossing to flies with black body-color (recessive), truncate behaves as a dominant, and manifests itself in some degree in nearly all of the F_1 flies that carry it. Two heterozygous factors, ordinarily recessive, may thus reenforce each other so that one becomes a dominant. Similar results were obtained in crosses with bar-eyed flies, and with star-eyed flies, but bar and star are both dominants themselves.

The dissociation of the truncate genotype

1. Truncate was crossed to stock having black body-color (chromosome II) and pink eye-color (chromosome III). In this cross, truncate could be used as a dominant (since black is present), and F_1 truncate males were back-crossed, separately, to black pink. In this way flies having all possible combinations of the chromosomes present in the truncate stock were obtained, and could be recognized, according to their sex, body color, and eye color, these latter being used as "identifying factors" for the chromosomes containing them. By then studying the amount of truncation in the flies having the different combinations, the effect of each of the chromosomes of the truncate stock, on the truncate character, could be determined. The following results were arrived at:

(a) There is a chief factor for truncate (T_2'), lying in the second chromosome; flies having only T_2', in heterozygous condition, and no other mutant factor for truncate, may occasionally have truncated wings. T_2' is partially dominant.

(b) The third chromosome of truncate stock also contains a mutant factor or factors for truncate, which may be designated as T_3'. This acts as an intensifier of T_2', and when it is heterozygous at least it produces no visible effect on the wing unless T_2' is present.

(c) These crosses, when made in both reciprocal forms, showed that there is a similar intensifier, T_1', in the first (X) chromosome, which intensifies truncate to about the same extent as does T_3'. When both intensifiers T_1' and T_3' are present they have a summative action. T_1', like T_3', produces no visible effect unless T_2' is present. T_1' is partially dominant, under the conditions of these experiments (with black present).

2. Crosses of truncate flies lacking T_1', obtained in these experiments, showed that the truncate character is also influenced by sex ("sex-limited"), in that it manifests itself more readily in females than in males.

3. Flies from these experiments, that were known to contain T_1' and T_3', but lacked T_2', were crossed with each other. As no wing modifications were produced, it was evident that both T_1' and T_3' together, in the absence of T_2', cannot cause truncation, or any visible effect on the wing, even when opportunity is given for them to become homozygous.

4. It was found possible to resynthesize truncate, according to calculation, from the flies of the non-truncate or imperfectly truncated classes of the dissociation experiment. Flies were chosen for mating together

which were scheduled, by their identifying factors, each to contain factors for truncate that were missing in the other. High-grade truncate then appeared in the offspring in the classes expected to contain it. This result furnished a confirmation of the conclusions from the dissociation experiment, and showed that the flies of the non-truncate classes had received the factors for truncate which they were supposed to have, and transmitted them regularly to their descendants.

5. In the crosses just mentioned, T_3' was sometimes introduced through the mother, sometimes through the father. The slightly lower grade of truncation, when T_3' is derived from the mother, indicates that T_3' is multiple in composition, consisting of two or more factors that may cross over from one another in the female.

6. Similar tests with T_2' indicated that this includes only one factor (at any rate one dominant factor).

7. It was found in later crosses that this "chief factor," T_2', is located in the "left-hand" end of the second chromosome, about 12 units to the right of star.

8. Experiments involving, in addition to black and pink, a fourth-chromosome factor, bent, showed that the truncate stock does not contain an intensifier in chromosome IV. In these latter experiments the distribution of the entire germ plasm was under observation.

Tests of the constancy of the factors for truncate

1. Individual tests were made of brother and of sister flies from the F_2 count of the "dissociation experiment," which could be seen, by their identifying factors, to have received from their father a given chromosome combination,—$T_1'\ T_2'\ T_3'$ in the case of the sisters, $T_2'\ T_3'$ in the case of the brothers. The tests showed that the differences in truncation between the flies having a given combination were not genetic; that is, the flies which receive the same chromosome combinations from their parents contain identical factors for truncate. It must be concluded from this that the factors for truncate follow exactly the distribution of the chromosomes, and that they do not undergo fluctuating variation.

2. A "pure-line" experiment was then undertaken, by repeated backcrossing of males of the above sort $(\frac{T_2'\ B}{b}\ \frac{T_3'\ P}{p})$ to black pink females. In each generation, in such a cross, males with the same truncate combination as their father are again produced, and may be recognized by their identifying characters (gray body and red eye-color). These are

then mated, as before, to black pink females from stock, and so the process may be repeated indefinitely. The danger of recombination is here excluded as effectually as in the case of self-fertilizing or asexually reproducing organisms, and so the purpose of a pure-line experiment is fulfilled. Two lines of flies were carried on in this way,—a "high" or plus-selected line, for twelve generations, and a "low" or minus line, for thirteen generations. The selection was absolutely ineffective; this corroborates the conclusion derived from tests of brothers and sisters, and shows that the factors for truncate are constant, and entirely contained in the (three large) chromosomes. It proves at the same time that the factors for truncate are not contaminated by their allelomorphs in the heterozygote.

The continual genetic variation occurring in the truncate stock must therefore be due to recombinations of factors for which the stock is perpetually heterozygous.

The cause of the inconstancy of the selected stock

1. To determine the reason for this perpetual heterozygosis of the stock, tests were undertaken to discover whether T_2' could exist homozygously. As it was next to impossible to determine this by direct tests, T_2' was placed in a chromosome with *b* (black), and flies with this combination were crossed to others having T_2' bound to *B* (gray). 72 offspring which had received at least one T_2' were then tested, by backcrossing to black, to discover whether they contained both the T_2' with black and the T_2' with gray. None of them, however, had more than one of the T_2' factors, and the ratios showed that T_2' acts as a lethal when homozygous.

Thus T_2' resembles the majority of dominant mutant factors of Drosophila in being lethal when homozygous, and pure stock of it cannot be obtained.

2. It follows that crosses of truncates should yield 1 non-viable: 2 potential truncates : 1 normal. As in the selected stock much fewer than one third of the flies that hatched were normal, tests were made to see whether any of the normals that should have occurred had been caused to appear truncate by reason of some other, recessive, "chief factor" for truncate. It was found, however, that no such factor existed; consequently the deficiency of normals was due, not to their appearing truncate, but to their actual absence,—this means that they were prevented from hatching by a lethal factor, which must have lain in the chromo-

some homologous to that bearing T_2'. Occasional crossovers between T_2' and the lethal would give rise to a few normals, however.

It is thus evident that there is in truncate stock a condition of "balanced lethal factors," similar to that existing in beaded Drosophila, Matthiola, and Oenothera.

3. Reciprocal out-crosses of truncate stock showed that T_1' was present, homozygous, in viable, fertile flies of the selected stock.

4. Crosses of flies from the dissociation experiment were arranged in such a way that T_3' was given opportunity to become homozygous, while T_1' and T_2' remained heterozygous. Many of the flies thus synthesized were as high-grade truncate as the best inbred stock, and much higher than the truncates from out-crosses. This proved that T_3', when homozygous, is not lethal, but that its effect on truncation is much stronger than when heterozygous. Females of this composition are so infertile, however, that it would be next to impossible to maintain stock pure for T_3'. This explains the occurrence of low-grade truncates, and normals carrying truncate, in the selected stock.

The origin of the truncate complex

1. Truncate is one of the most frequent characters to appear sporadically, by apparent mutation, in various stocks of Drosophila. In most of these cases, however, intensifiers of truncate are absent, and so the possibility is not excluded that truncate was present previously in the stocks, and merely failed to manifest itself before. In accordance with these circumstances it is usually found that the truncate fails to reappear in the descendants of such flies, except in a minute percentage of individuals.

When, however, truncate appears in stocks containing bar, or other factors which themselves intensify truncate, it may be concluded that truncate has here arisen as a real mutation. Several such truncates have been tested, and found to be inherited, and evidence was obtained that their mutant factor for truncate was identical with T_2', the chief factor of ordinary truncate stock. None of the intensifiers of the ordinary stock were present, however.

2. The hereditary behavior of the original truncate shows that one of the factors for truncate had been present, undetected, in the parental long-winged stock, and that the appearance of truncate was due to the origin, by mutation, of a second factor. Thus the first truncate-winged fly contained T_2' and an intensifier. The other mutant factors of the

truncate stock appeared later, and they were perpetuated through the process of selection, because they influenced the truncation in the direction desired.

It is of significance for evolution, as well as for genetics, that particular races may thus have pre-existing in them factors which by themselves are invisible, but which favor the manifestation of a certain character.

3. The occurrence of more modifiers for truncate than for most characters is explained not only by the fact that the modifiers were specially picked out by selection in the case of truncate, but also by the marked instability of the developmental reaction whereby the truncate character is produced. This instability, or susceptibility of modification, is disclosed by (1) the somatic variability of the character, when the genotype is kept constant, and (2) by the number of mutant factors for other characters which act upon truncate in addition. On account of this modifiability it should happen that relatively many of the mutant genes which arise would be able to affect truncate, and it should therefore be easier to find either intensifiers or "inhibitors" for truncate than for most characters. The reason for the readier discovery of modifiers would hence be, not that the process of mutation is influenced in some way by truncate, but, on the contrary, that truncate is especially susceptible of being influenced by mutation.

The existence, in certain races, of factors like T_2', which make a certain character (here, the wing shape) more highly modifiable, may be of particular importance in evolution.

General applications

1. The results described in the first section of this resumé would undoubtedly have been regarded by certain writers as positive proof of the ineradicable fluctuability, miscibility, or, as it were, fluidity, of a unit character. The case was, in fact, more extreme than those upon which they rely as evidence for their doctrine. And yet it has been found that the inheritance of this character is entirely confined to the recognized chromosome system, and depends on definite chromosomal factors which are neither miscible nor inconstant.

Although mutations affecting this character may occur with somewhat greater frequency than those for many other characters, the mutations are not nearly of the order of frequency which would be necessary in order for them, by themselves, to give rise directly to a true genetic fluctuability. Practically all the genetic variability and modifiability of

the stock is, on the contrary, due to the occurrence of recombinations among those mutant factors which have, on rare occasions, previously arisen, and variation due to a real new mutation, although possible, is highly exceptional. Even in such cases, moreover, the mutations do not all consist in variations of some one particular factor, but they may affect any one of many factors that are concerned with the character.

The use of cases resembling the truncate case, in order to support the doctrine of fluctuating variability of single unit-factors, is therefore entirely unjustifiable, in the absence of analyses comparable with those made in this case.

2. It is believed that the general methods developed in this case will become increasingly useful with the growth of knowledge of the linkage groups in organisms. It has been shown how, by the use of "identifying factors," any given genotype may be cut up for study, put together again in various ways, or held in a particular combination for the maintenance of stock or for observations on factor constancy. At the same time the distribution among the offspring, of the entire (chromosomal) germ plasm of a parent may be completely exposed to view.

The suggestion is made that such methods may also be necessary before much progress can be made in the study of the more important characters in human genetics.

LITERATURE CITED

DUNCAN, F. N., 1915 An attempt to produce mutations through hybridization. Amer. Nat. **49**: 575-582.

HYDE, R. R., 1914 Fertility and sterility in *Drosophila ampelophila* I-IV. Jour. Exp. Zoöl. **17**: 140-212, 343-372.

MORGAN, T. H., STURTEVANT, A. H., MULLER, H. J., and BRIDGES, C. B., 1915 The mechanism of Mendelian heredity. 262 pp. New York: Henry Holt and Co.

MULLER, H. J., 1917 An Oenothera-like case in Drosophila. Proc. Nation. Acad. Sci **3**: 619-626.

1918 Genetic variability, twin hybrids and constant hybrids in a case of balanced lethal factors. Genetics **3**: 422-499.

SPOTTING INHERITANCE IN *DROSOPHILA BUSCKII* COQ.[1]

DON C. WARREN

Georgia State Board of Entomology, Valdosta, Georgia

[Received July 26, 1919]

TABLE OF CONTENTS

INTRODUCTION 61
The problem 61
Description of species 61
Material and methods of breeding 62
Male spotting 63
Selection lines 63
Selection in line 500 64
Results 67
Selection line 502 68
Results 70
Selection line 504 71
Results 73
Discussion of selection experiments 74
Permanency of the strains established by selection 75
Discussion of behavior of male line 501 75
Genetic behavior of strain 501 when crossed to strain 504 low 78
Effect of temperature on the male spotting 81
Crosses between the various male strains 84
Summary 85
Female spotting 86
Abnormal female 86
Selection of ab strain 86
Matings of ab strain with normal strain 500 low 91
Summary of various types of mating in testing the behavior of ab when crossed to the strain 500 low 96
Crosses between the ab strain and strain 502 high 96
Genetic behavior of ab strain when crossed to various strains 98
Relation of temperature to genetic behavior of abnormal female condition 99
Linkage 101
Attempts to establish an ab strain from other stocks 102
Discussion of the analysis of the abnormal type of female spotting 104
Summary 105
GENERAL DISCUSSION 106
GENERAL SUMMARY 108
LITERATURE CITED 109

[1] From the Zoölogical Laboratory of INDIANA UNIVERSITY. Contribution No. 171.

INTRODUCTION

Few groups of animals or plants have received so much study by the geneticist as the fruit fly of the genus Drosophila. This work, carried on largely by MORGAN and his co-workers, has consisted almost entirely of an analysis of the inter-relations of variations found in the laboratory. Large numbers of flies have been minutely examined for germinal variations and this critical study has resulted in the finding of a large number of mutations of various sorts. These mutations were usually observed soon after their appearance and further work with them has resulted in adding much new and valuable light to our knowledge of heredity.

Although many mutations have been found in the laboratory, very few have been recorded in flies taken from nature. This fact has led some writers to question whether these variations found in the laboratory are really potent factors in the evolutionary process. The difference may lie in the fact that flies in nature have not been given the exhaustive study which has revealed the mutations appearing in the laboratory. However, when one examines any large number of flies in nature he is struck, not by the large number of variations which may be of evolutionary value, but rather by the lack of these. There is certainly very little evidence of any variations similar to those found in the laboratory, which seem to be taking the place of any existing contrasted characters.

THE PROBLEM

The problem of this paper has been to analyze variations which were found in nature. In studies of this sort it is not possible to know anything concerning the origin of the variations and hence the analysis is more difficult. The variation studied was the spotting of the abdomen. The study of the variations in the male, together with a variation in female spotting, which later appeared in the writer's stocks, composes the work of this paper. In its final form the problem became an attempt to analyze critically the factors which govern or influence the spotting character or characters in the fly. To narrow the field still further the writer has limited himself to the spots on a single segment of the abdomen. Whether the inheritance of the spots on the other segments of the abdomen are dependent upon the same factors remains in question.

DESCRIPTION OF THE SPECIES

The fruit fly *Drosophila busckii* is slightly smaller than the well known

Drosophila melanogaster and is readily distinguished from it by conspicuous spots on the abdominal segments. Each segment, excluding the last two, bears six spots. These spots are arranged on each segment as follows. There is a pair of rather large distinct spots on the dorsal surface of the segment, one on each side of a mid-dorsal line. A second pair appears more laterally, one on each side just below the last-mentioned pair. The third pair of spots are small ones, latero-ventral in position. The spots are arranged so as to make six rows running lengthwise of the abdomen. There is seldom any variation in the spotting of the above-described segments.

The spotting of the fifth segment differs from these, however, and the variations found on that segment furnish material for this study. In the female it is like that of the other more anterior abdominal segments,[2] but in the male there are variations. The large dorsal pair of spots is always present and the presence of this pair alone seems to be the more common type of spotting among males found in nature. Either one or both of the other two pairs may be present or absent or any one spot of a pair may be combined with any other spots, thus giving all the possible combinations of the constant dorsal pair and the other four variable spots. These various combinations are shown in figure 1. The spots show a tendency to be present or absent in pairs, that is, if the outside spot on one side is missing the corresponding spot on the other side is also likely to be missing. This tendency is especially true of the outside pair of spots and one outside spot occurs without the other so infrequently that no record of this type of spotting has been made in the tables which follow.

MATERIAL AND METHODS OF BREEDING

The flies used in these experiments were collected in the vicinity of Bloomington, Indiana, during the autumns of 1915 and 1916. In nature they breed in decaying fruit and vegetable matter but most frequently in rotting tomatoes. At first it was attempted to breed them on banana in the laboratory. It was found that they could be reared on this food, but with some difficulty. Later, moistened bran was tried and used very successfully, being a food which is both convenient to handle and one upon which the *Drosophila busckii* breeds very well. This species is very prolific and requires about fourteen to twenty-five days for a generation, depending upon the temperature. It does not well withstand high temperatures and the extreme summer heat encountered in central Indiana sometimes killed stocks of flies.

[2] One exception was found which will be discussed later in the paper.



FIGURE 1.—Diagram of the various types of spotting found on the fifth abdominal segment of the male. Each type represents a flattened segment showing the spots upon it. The large dorsal pair was always present and was found in every possible combination with the other four smaller spots. The spots tended to be present or absent in pairs and are referred to throughout the paper as the dorsal pair, the middle pair and outside pair. Type 1 was most common in nature.

MALE SPOTTING

Selection lines

This study was suggested by the observation that there were variations in the spotting of the males. Preliminary work upon these variations indicated that they were heritable to a certain degree. The earlier work consisted largely of crosses between the more extreme variations to see how they would behave, genetically. Further than establishing to the writer's satisfaction that these variations were heritable, very little was accomplished, for no pure strains of any of the variations had been acquired. So it was decided that the best method of procedure would be to try to establish by selection, strains which were pure for some of these variations. Selection was made for the two extremes of the various types of spotting in the males. These two extremes were the normal or two-spotted condition (fig. 1, type 1, and plate 1, no. 1) and the six-spotted condition (see fig. 1, type 16, and plate 1, no. 3). As well as


1
2
3
4
5
6
7
8

LEGEND FOR PLATE I

Some of the types of spotting found in *Drosophila busckii*. The work of this paper is confined to the fifth abdominal segment (the last segment showing spots).

No. 1.—Normal male with only the dorsal spots.
No. 2.—Male with dorsal spots and small outside one.
No. 3.—Male having full set of three pairs of spots.
No. 4.—Normal female with full set of three pairs of spots.
No. 5.—Abnormal female with middle pair of spots missing.
No. 6.—501 type of male with very large outside spots and also small middle spot.
No. 7.—Male with dorsal spots and small middle spot.
No. 8.—501 type of male with large outside spot.

being an attempt to isolate two desired types of males the problem was outlined as a selection problem. From flies taken in nature three stocks were started which were not known to be related. To begin with quite a number were examined in each stock to learn what was the ratio of the various types of male spotting in each. Then from among the offspring of a single pair in each stock a "high" and a "low" strain were started. The high strain was one in which the paternal parent in each succeeding generation was to be a male with a full set of six spots, while the low strain was one in which the father of each generation was to be a male of the normal or two-spotted type. At the same time that these two strains were started a control strain was also started in each stock. In the control the parents were selected at random with no knowledge of the nature of the spotting of the fathers. The three stocks in which the selection was started were known as 500, 502 and 504 and after the selection was started there was a low strain, a high strain and a control strain in each.

Selection line 500

Three pairs of flies were taken in nature and placed in a bottle which was numbered 500. This was the origin of the strain 500. A record was made of the spotting of the offspring from this bottle and six pairs of the offspring were mated *inter se*.[3] Of these six matings, one was used as the starting point of the high and low strains of line 500. This mating was a single-pair mating and the female was virgin, so all of the offspring were necessarily full brothers and sister. This gave the two strains as nearly uniform starting conditions as possible. The only difference in conditions at the starting of the high and low strains was that two-spotted males were used in the low strain and six-spotted males were used in the high strain. In as far as possible uniformity of all conditions except the one mentioned was maintained during succeeding generations of selection. Tables 1, 2, and 3 show the selection in the three strains of line 500.

In this line a control was not started at the beginning of selection but was originated by mating the high and low strains at generation 3. All succeeding generations of the control were mated at random with no knowledge of the type of male used.

[3] It will be noted that the counts obtained from the earlier matings were rather small as compared with the later generations of selection. This was due to the fact that the flies were at first reared on banana upon which they do not thrive well. Later wet bran was used and found much more satisfactory.

TABLE 1
Selection in strain 500 low.

Generation	Percent selected type	Types of male spotting										Total males
		Type 1	Type 2	Type 4	Type 5	Type 6	Type 11	Type 12	Type 13	Type 15	Type 16	
Counts before selection	56	152	14	11	7	7	22		9	3	48	273
1	28	91	9	13	5	3	145	7	6	11	36	326
2	85	430	4		2		66		2		1	505
3	92	1166	6	11	12	1	46	1	8		17	1268
4	96	606	7	6	6	1	8	1	2		5	642
5	99	717	4		4							725
6	94	239	5		5	2		1	1		1	254
7	97	477	3		10	1			1		1	493
8	97	433	3		10							446
9	99	347	2		1							350
10	94	546	5	2	6	8	9				3	579
11	93	435	7		5	2						469
12	99	372	1		1							374
13	99	350	1		1		1				1	354
15	89	181	1		1	1	17	1			2	204

Table 1 shows the selection for the two-spotted type of male (type 1) in strain 500 low. The column showing the percent of selected type gives a fair idea of what was accomplished by selection. The column headings, type 1, type 2, etc., refer to the types of male spotting found. These types are shown in figure 1. Some of the types given in figure 1 are omitted in the tables which follow. This is due to the fact that there were so few individuals in some of the types that space was not given them in the tables.

In generation one of the low strain eight matings were made but in practically all of the succeeding generations two stock matings were made in each generation of each strain. The graph shown in figure 2 gives a fair idea of the results obtained by selection in this line. Selection was much more effective in the direction of the two-spotted males. The graph representing the low strain in generation 4 drops to near one-hundred percent two-spotted males and remains so during the rest of the selection. Selection in the opposite direction was less progressive. The graph representing the high strain is very irregular and a high percentage of six-spotted males was not reached till generation 9. During the first eight generations of selection the high strain frequently threw more two-spotted males than six-spotted ones. This seemed to be characteristic of line 500. There was a tendency in all strains to produce a high number of two-spotted males and the percentage of six-spotted was difficult to increase. The control in which there was no selection, al-

TABLE 2
Selection in strain 500 high.

Generation	Percent selected type	Types of male spotting										Total males
		Type 1	Type 2	Type 4	Type 5	Type 6	Type 11	Type 12	Type 13	Type 15	Type 16	
Counts before selection	18	152	14	11	7	7	22		9	3	48	273
1	21	10		1			17	1		2	8	39
2	43	29	1	4	3	1	22	4	2	2	50	118
3	24	179	8	24	7	4	47	4	24	7	98	402
4	43	122	6	27	7	4	51	5	22	6	186	436
5	7	348	19	15	25	22			17		35	481
6	12	360	20	22	26	22	5	4	12	3	62	536
7	17	335	19	28	20	18	27	4	26	6	100	583
8	34	146	2	18	14	7	90	19	18	15	171	500
9	79	1		2		1	13	10	2	8	138	175
10	61	99	10	12	23	28	49	23	6	24	422	696
11	56	22	1	3	2		134	21	2	18	261	464
12	86	9	3	2	6	4	13	4		2	268	311
14	66	8					42	5		3	111	169

Table 2 shows the selection for the six-spotted type of male (type 16) in strain 500 high. The types referred to in the column headings are found in figure 1. The increase in percent observed as one reads down the second column shows what was accomplished by selection.

TABLE 3
Counts for control strain of line 500.

Generation	Types of male spotting										Total males
	Type 1	Type 2	Type 4	Type 5	Type 6	Type 11	Type 12	Type 13	Type 15	Type 16	
Counts before selection	152	14	11	7	7	22		9	3	48	273
1	195	6	1	6		11	2	1	1	11	234
2	134	1		1	1						137
3	184	5	5	12	2		1	2		6	217
4	209	6	2	5	3			1			226
5	199	8	8	4	8	45	3	3	6	23	307
6	179	2		2	1						184
7	123	1		4	1						129
8	109	3	1	4	3	9				10	139
9	148	13	2	10	6	6	1		1	1	188

Table 3 shows the counts for each generation of the unselected strain 500 control. These counts should be compared with those in tables 7 and 2. The behavior of this strain in relation to the two selected strains in line 500 is best shown in figure 2.



FIGURE 2.—Results of selection in line 500. The dash line represents the high strain, the dotted line the control, the solid line the low strain. The abscissae represent the generations of selection while the ordinates represent the difference between the percentage of two-spotted and six-spotted males in any generation. Differences in favor of the six-spotted males are plotted above the zero line while those in favor of the two-spotted males are plotted below the line; *e.g.*, in generation 2 of the high strain, 43 percent of the males were six-spotted and 24 percent were two-spotted, so the difference was 19 in favor of the six-spotted males and was plotted at 19 above the zero line. The data for this graph are found in tables 1, 2 and 3.

though somewhat more irregular, did not range far from the low strain. The extra spots of the six-spotted males of the high strain were all rather indistinct as compared with those in line 502.

Results

There can be no doubt but that the high and low strains have been made different by selection. Although a much lower percentage of males of the desired type were obtained in the high strain than in the low strain, it is evident that more was really accomplished by selection

in the high strain. This is true because the line 500 at the beginning had a tendency to produce two-spotted males and by selection a greater difference was brought about between the high strain and the control than between the control and the low strain.

Selection line 502

This selection experiment (tables 4, 5 and 6) was started and carried out in the same manner as on line 500. The initial stock was taken from nature, bred in the laboratory for two generations and then from among the offspring of a single pair the selection strains and control were started. This line soon showed itself to be characteristically different from line 500. Line 502 showed a tendency to produce a high percentage of six-spotted males. The graph (figure 3) representing the behavior of the low strain in this line was much more irregular than the

TABLE 4
Selection in strain 502 low.

Generation	Percent selected type	Types of male spotting										Total males
		Type 1	Type 2	Type 4	Type 5	Type 6	Type 11	Type 12	Type 13	Type 15	Type 16	
Counts before selection	31	78	8	14	4	10	30	1	10		98	253
1	22	26		1	1		73	2		3	11	117
2	24	96		9	2	3	123	6	12	8	135	394
3	82	173	1	2	2		26		2		4	210
4	94	739	5	14	7	1	12		8		2	788
5	85	622	24	3	26	9	4	4	7		28	727
6	84	722	19	7	23	15	14	3	11	2	39	855
7	93	480	7	7	13	4	2		1	1	3	518
8	94	410	4	1	4	2		1			2	424
9	79	586	18	13	25	23	33	5	6	8	27	744
10	32	92		1	4	1	166	7	1	4	8	284
11	54	230	3	2	6	5	139	8	5	15	12	425
12	81	370	8	3	10	1	52	4		3	7	458
13	50	78	3	1		1	48	13		2	9	155
15	47	98	2		1		94	1		2	11	209

The counts in table 4 show the results of selection for the two-spotted type of male (type 1) in strain 502. The types referred to in the column headings are shown in figure 1. The second column shows the effect of selection and the decrease in the percent of selected type in the latter part of the experiment will be seen to be due to an increase in the type 11 males. The type 11 male (see figure 1) is one with only the outside spots and these spots are affected by temperature variations. Temperature is probably the cause of the decreased percent in the last few generations.

TABLE 5
Selection in strain 502 high.

Generation	Percent selected type	Types of male spotting										Total males
		Type 1	Type 2	Type 4	Type 5	Type 6	Type 11	Type 12	Type 13	Type 15	Type 16	
Counts before selection	38	78	8	14	4	10	30	1	10		98	253
1	11	8					44	3		2	7	64
2	62	48	1	7	3		108	4	7	8	305	491
3	70	33		8	2	1	40	5	8	10	246	353
4	60	26	1	9			127	28	6	18	326	541
5	75	33	2	7	3	2	62	13	5	9	416	552
6	70	2	1	4			104	20	2	20	351	504
7	89			1			38	20	1	16	585	661
8	55			1			141	31		7	217	397
9	86						10	2		4	96	112
10	100									1	502	503
11	93						24	1			351	376
12	100						1	1			458	460
14	98						1			2	147	150

Table 5 gives the results of selection in strain 502 high. Selection was for the six-spotted or type-16 male. Figure 1 gives a diagram of the types referred to in this table. It will be seen from column two that it was possible by selection to obtain almost one hundred percent of the desired type.

TABLE 6
Counts for control strain of line 502.

Generation	Types of male spotting										Total males
	Type 1	Type 2	Type 4	Type 5	Type 6	Type 11	Type 12	Type 13	Type 15	Type 16	
Counts before selection	78	8	14	4	10	30	1	10		98	253
3	374	3	7	6	1	83	4	9	3	16	506
4	166	5	2	10	9	6	2		1	21	222
5	89	3			2			1		1	96
6	135	5	3	5	5	13	3	3	3	19	194
7	92	2	7	5	4	35	9	5	5	88	252
8	38		4	2	3	33	7	1	5	35	128
9	30	2	2	1		26		1	1	17	80
10	7		1			98	10		1	66	183
11	62		1	8	6	33	19		2	44	175

The counts in table 6 represent the behavior of strain 502 control. This was the unselected strain of line 502. The best comparison of the behavior of this strain with the high and low strains of 502 is shown in the graph in figure 3.



FIGURE 3.—Results of selection in line 502. See figure 2 for explanation of graph. The dash line represents the high strain, the dotted line the control and the solid line the low strain. Tables 4, 5 and 6 give the data for this graph.

corresponding one in line 500. Also the percentage of two-spotted males in the corresponding generations was lower than in strain 500 low. The percentage of six-spotted males in the high strain of 502 gradually increases and in the last few generations nearly approaches one hundred. The apparent loss of effect of selection in the last few generations in the low strain was probably due to temperature and will be discussed later. The extra spots on the males of the high strain of 502 were very distinct.

Results

Here again there is clear-cut evidence of results having been accomplished by selection. The graphs representing the two strains show a wide divergence and the control, although rather irregular, occupies an approximately midway position. There was a greater difference brought about by selection between both strains and the control than in line 500.

Selection line 504

The selection strains in this experiment were started as in the other two experiments. The data for this line are found in tables 7, 8 and 9. This line behaved somewhat similarly to line 500 and perhaps a little more extremely. There was a tendency to produce two-spotted males. Very early in selection (generation 2) the low strain produced nearly one-hundred percent two-spotted males as is shown by the graph and data for this strain. The graph (figure 4) then runs very uniformly near one-hundred percent except in the last two generations when there was a rather sudden rise. This irregularity was due to temperature and will be discussed under the topic of temperature. In the high strain the results were rather variable for the first few generations of selection. The percentage of six-spotted males was hard to increase and not until the last few generations of selection did the graph representing the behavior of the high strain rise above the zero line, thus showing a preponderance

TABLE 7

Selection in strain 504 low.

Generation	Percent selected type	Types of male spotting										Total males
		Type 1	Type 2	Type 4	Type 5	Type 6	Type 11	Type 12	Type 13	Type 15	Type 16	
Counts before selection	31	117	15	14	7	14	89	7	12	1	97	373
1	93	332	1				25					358
2	98	1287	4	1	4		9	1	2		1	1309
3	99	632	1		1		1					635
4	100	235										235
5	98	437	2		2	4						445
6	99	547	5		2							554
7	99	458	2		1				1			462
8	98	452	4	1	4	1						462
9	92	462	3	1	3	1	29		1	1		501
10	97	302	2		5	1						310
11	99	389	1									390
12	98	406		1	3	1	3					414
14	62	133	1		1		78	2				215

Table 7 gives the selection for the two-spotted male (type 1) in strain 504 low. The types referred to in this table are shown in figure 1. It is seen that selection was very effective in this strain. All of the strains of line 504 showed a tendency to produce males of the two-spotted type. In this respect this strain is quite different from the low strain of line 502. By comparing the results of this table with those in table 4 this difference is readily seen.

TABLE 8

Selection in strain 504 high.

Generation	Percent selected type	Types of male spotting										Total males
		Type 1	Type 2	Type 4	Type 5	Type 6	Type 11	Type 12	Type 13	Type 15	Type 16	
Counts before selection	26	117	15	14	7	14	89	7	12	1	97	373
1	2	106	4		2		81	2			4	199
2	3	384	6	3	6		45	2	5	1	14	466
3	4	728	27	13	42	33	67	3	16	3	34	966
4	3	521	17	5	16	12	50	4	6	3	21	655
5	10	429	14	13	17	15	193	23	8	13	80	805
6	6	429	18	19	24	26	117	21	8	9	46	717
7	6	219	12	6	20	12	89	8	7	3	22	398
8	14	238	19	14	17	15	170	19	5	20	82	599
9	4	39	2		1		199	6	2	13	10	272
10	32	2		1		1	221	25		18	127	395
11	51	62	8	6	19	14	86	16	5	8	230	454

In table 8 are shown the results of selection in strain 504 high. It was difficult by selection to increase the percent of six-spotted males (type 16) due to the tendency of all of the strains of this line to produce two-spotted males. However by examining the graph in figure 4 it will be seen that selection made the two selected strains act quite differently.

TABLE 9

Counts for control strain of line 504.

Generation	Types of male spotting										Total males
	Type 1	Type 2	Type 4	Type 5	Type 6	Type 11	Type 12	Type 13	Type 15	Type 16	
Counts before selection	117	15	14	7	14	89	7	12	1	97	373
1	290	7	1	15	1	33	1	3	1	8	360
2	131	1		4							136
3	138	3		1	2	8	1	1	1	2	157
4	193	5	1	7	5			1			212
5	242	8	4	7	3	18					282
6	105	1			1						107
8	147	7		7	1					5	167
9	114	2		7	1			1			125
11	110	1	1	4	2	11					129

Table 9 shows the counts for each generation of the unselected strain 504 control. The behavior of this strain in its relation to the two selected strains of line 504 is shown in figure 4.



FIGURE 4.—Results of selection in line 504. See figure 2 for explanation of graph. The dash line represents the high strain, the dotted line the control, and the solid line the low strain. Tables 7, 8 and 9 give the data for this graph.

of six-spotted males. The percentage of six-spotted males never rose as high as in the high strains of lines 500 and 502. The extra spots on the six-spotted males in the high strain were much less distinct than in line 502. The control was not started from the original stock of line 504 but was originated as a cross between the high and low strains after two selections. The control threw a rather high percentage of two-spotted males but is shown by the graph as running more irregularly and slightly above the low strain.

Results

Although in line 504 it was impossible by selection to produce a very high percentage of six-spotted males, there can be no question about the significance of the results. The divergence of the two lines repre-

senting the high and low strains clearly shows what has been accomplished.

Discussion of the selection experiments

Selection has been carried out upon three distinct stocks of flies. In each case two strains have been produced which behave quite differently with respect to the character upon which selection acts. As to how selection acted is a more difficult problem. There are two conflicting views: first, the pure-line view that selection can only isolate existing pure lines, or the elimination and accumulation of multiple factors; second, the opposing view until recently held by CASTLE that selection modifies the gene.

This experiment offers no demonstration of the correctness of either view. In the experiment under consideration the object of selection was not to increase and decrease the intensity of a given character but rather to vary the percentage of individuals showing the character. The character was the presence or absence of four spots on a given abdominal segment. It was found that each line selected from nature seemed to have a more or less characteristic behavior with respect to the presence or absence of these spots. The ease with which a strain could be varied by selection in a given direction and the range of variation seemed to be a characteristic of the strain itself. The easiest explanation of this behavior is that this is a case of isolating pure lines. This in a way would be comparable to the classic experiments of JOHANNSEN upon beans. The difference in range of variation of the different strains is hardly sufficiently great to make a good case of this sort. However the rather striking difference in amenability to selection in a given direction does lend support to the pure-line theory and causes the writer to believe that the lines 500, 502, and 504 were all different genetically when taken from nature. It should be added, however, that at the end of the selection experiment the two high strains of lines 502 and 500 (see discussion of results of crosses between various male strains, which follow later in the paper) were crossed and the F_1 results indicated that at least some of the factors for spotting were common to both high strains. So although the two high strains reacted differently to selection and even remained somewhat different at the end of selection, they appeared to have the same genetic constitution at the end. Selection may have changed the genetic constitution of one of them. The writer is unable to see how the assumption of multiple factors would assist in explanation of this case and therefore merely presents the observed facts allowing the reader to give his own interpretation.

Permanency of the strains established by selection

An experiment was carried out to test whether the difference brought about by selection in the high and low strains was permanent. Line 502 was used in this test and at generation 5 of selection, a strain was isolated from each of the high and low strains. These strains were to be carried along with the selection strains but the male parents of each generation were to be taken at random with no knowledge of the type of spotting which they possessed. Random selection was carried on for seven generations and then counts were taken of the offspring. The results (table 10) show that there had been no retrogression in either strain in spite of the fact that selection had been stopped for several generations. In the high strain the count showed that seventy-six percent of the males were of the six-spotted type and at generation 5 when this strain was isolated to be carried without selection, the strain showed seventy-five percent of the males to be of the six-spotted type. So the high strain had not changed. At the time that selection was stopped, the low strain showed eighty-five percent of the males to be two-spotted and after seven generations of mass culture without selection it still showed eighty-nine percent of the males to be two-spotted. These two strains were carried for twenty-three generations without selection and the intention was to make another count after a long period. Pressure of other work prevented these counts from being made and finally the two strains were killed by the extreme summer heat of Alabama where the writer was then located. However, in transferring the flies from generation to generation it was observed that there always remained a striking difference in the spotting of the two strains and it is believed that the effects of selection remained permanent.

Discussion of behavior of male line 501

Before discussing the genetic behavior of the various selection strains it will be well to consider an unusual type of male spotting which was found in the stocks carried. A number of stocks of flies were taken from nature at the time that the male selection lines were started. One of these stocks (line 501) was observed to produce males which had unusually large outside spots on the fifth abdominal segment (see plate 1, no. 8, page 64). The spots of this outside pair were frequently almost as large as those of the dorsal pair. This was unusual, for these spots were ordinarily much smaller than the dorsal ones. Selection was started using the most extreme males in an attempt to establish a pure

TABLE 10
Permanency of effect of selection.

A. After five generations of selection

Strain	Percent selected type	Type 1	Type 2	Type 4	Type 5	Type 6	Type 11	Type 12	Type 13	Type 15	Type 16	Total males
502 low	85	622	24	3	26	9	4	4	7		28	727
502 high	75	33	2	7	3	2	62	13	5	9	416	552

B. Same strain after seven additional generations without selection

502 low	89	281	2		1		32	1				317
502 high	76	4			1		45	13		4	217	284

Table 10 shows the permanency of the effect of selection in line 502. Strain 502 low and 502 high originally came from the same stock. In strain 502 low selection was made for type 1 males while in strain 502 high selection was for type 16 males. (These types are shown in figure 1.) After 5 generations of selection in which considerable difference was brought about, stocks were taken from the two strains and carried unselected for seven generations. At the end it is seen that the difference in the two strains remained permanent.

stock of this type of male. Immediately an almost pure stock was obtained. Practically all of the males possessed the outside pair of spots and these spots were unusually large in most cases. Some of the males also possessed the middle pair of spots (plate 1, no. 6) making a male of the six-spotted type. Selection was carried on for several generations until practically all of the males possessed the unusual outside spots.

Genetic behavior of line 501 when crossed to strain 504 low

The 501 line was crossed with the low strain of 504. It will be remembered that the low strain of 504 was practically pure for the two-spotted condition or the absence of the outside pair of spots which were characteristic of line 501. The 501 males and females were each crossed with the opposite sex from 504 low. The F_1 results are shown in table 11. Eleven matings were made by crossing the 504 females, by the 501 males, and ten matings of the reciprocal cross. The data for these crosses show a rather striking difference in the results where the 504 female was crossed with a 501 male, as compared with the cross in the opposite direction. When the female was from 501 there was a much higher percentage of the males in the F_1 showing one or both spots of the middle pair than when the male of the same line was used. This is not a chance fluctuation but holds true for practically every mating. The

middle pair of spots was not the one which was of primary interest in this set of matings for only a small percentage of the males of 501 possess the middle spots. From figure 1 it can be seen that all of the types listed in table 11, except types 1 and 11, possess one or both of the middle spots. When the 501 female was used in the cross, about thirty percent of the males in the F_1 have one or both of the middle spots, but when the 501 male was used, less than three percent of the males possessed one or both of the middle spots. The only explanation that can be offered for this peculiarity is that one or more of the genes affecting the presence of the middle spots are located in the sex chromosome. Not all

TABLE 11

Strain 501 × strain 504 low.

F_1 generation

Type of mating	Type 1	Type 2	Type 4	Type 5	Type 6	Type 11	Type 12	Type 13	Type 15	Type 16	Total males
504 ♀ × 501 ♂	373		1			1539	18	2	13	18	1964
501 ♀ × 504 ♂	244	4	8	2	1	953	90	8	105	311	1726

F_2 generation

Type of mating	Type 1	Type 2	Type 4	Type 5	Type 6	Type faint 11	Type 11	Type 12	Type 13	Type 15	Type 16	Total males
Faint type 11 ♂ from cross 504 ♀ × 501 ♂	314	14	10	24	19	613	398	42	2	27	149	1612
Type 16 ♂ from cross 501 ♀ × 504 ♂	357	19	1	17	15	694	433	38	1	20	148	1743
Good type 11 ♂ from cross 501 ♀ × 504 ♂	376	13	3	9	11	667	416	26	2	14	95	1632
Total for F_2 generation	1047	46	14	50	45	1974	1247	106	5	61	392	4987

Table 11 shows the results of the cross between strain 501 and strain 504 low. Strain 501 possessed very large outside spots while strain 504 lacked them. In the F_1 results note the difference in percent of type 16 males depending on whether the male or female from 501 was used in the cross. The F_2 results indicate that it makes no difference in the results of that generation what sort of an F_1 male is used as father of the F_2 generation.

of the genes for the middle spots can be in the sex chromosome, for in that case a female from 501 should transmit the character to all of her sons in the F_1 generation. After observing this tendency toward sex-linked inheritance of the middle pair of spots in 501 the writer examined some of the other data in which the middle pair of spots was involved tc

see if the same behavior had been overlooked there. A cross had been made between a strain known as the abnormal strain[4] and 502 high. In the abnormal strain the males were practically all two-spotted or lacking both in the middle and outside pair of spots, while the males of the 502 high possessed all three pairs. In the crosses between these two stocks, there was no difference in the F_1 results whether the male or female from 502 high was used. There is no indication of sex-linked inheritance with respect to the middle pair of spots, though such linkage should have been manifest in this cross since the abnormal strain was practically pure for the absence of the middle pair of spots while the 502 high was practically pure for their presence. Thus it seems necessary to conclude that there is a gene in the sex chromosome in line 501 which influences the presence of the middle pair of spots while in strain 502 high none of the genes influencing the inheritance of the middle pair of spots are in the sex-chromosome.

Turning now to the genetic behavior of the outside pair of spots for which the cross between 501 and 504 low was made, it will be seen from table 11 that in the F_1 there is a more or less incomplete dominance of the presence of the outside spots in 501 over their absence in 504 low. The count shown in the first column (type 1) of the table represents the males with neither outside spot while practically all of the other columns were for males which possessed one or both of the outside spots. As to the percentage of the males in the F_1 showing the outside spots, it seems to have made no difference whether the male or female from 501 was used in the cross. However, one difference was evident in the F_1 generation between the crosses where the 501 male or female was used in the mating. When the female of the cross was from 501 the male offspring showing the outside spots in the F_1 showed them in greater intensity than when the 501 male was used in the cross. In the cross of 501 male by 504 female, the outside spots which appeared on the males of the F_1 generation were rather faint. When the cross was made in the opposite direction many of the male offspring were almost typical 501 males with respect to the intensity of the spots. The F_2 results are also given in table 11. Three types of matings were made in the F_1 generation. From the F_1 of the cross 501 female by 504 male, both the six-spotted males and the almost typical 501 males were mated to sisters while in the F_1 of the cross 504 female by 501 male, the males showing the outside spots faintly were mated to sisters. The F_2 results show no difference between these three sorts of matings. In the table of the F_2 results the

[4] The behavior of this strain is discussed later in this paper.

column headed type 11 represents the count of males which were typical 501 while the column just preceding it contains males with the outside spots showing faintly. The males which show the outside spots faintly are probably heterozygous while the apparently typical 501 males are probably pure, segregated males of that type. The males in the first column are typical 504 males with only the dorsal pair of spots. Some of these typical 501 and 504 males were taken from the F_2 generation and mated back to females from their respective lines. The resulting offspring were typical of the line to which the males were backcrossed showing that the two types of males used in the cross had segregated after the Mendelian manner.

These crosses indicate that the outside pair of spots which are characteristic of line 501 is inherited in the usual Mendelian manner except that there is a lack of dominance of its presence or absence. There is a difference in the intensity of the outside spots on males of the F_1 generation depending upon whether the 501 male or female is used in the cross. The more intense spots appear when the 501 female is used. This might be taken to indicate that a sex-linked factor is involved in the inheritance of the outside spots. There is fairly definite evidence from the crosses made that sex-linked inheritance is involved in the transmission of the middle pair of spots in 501 although there is no evidence that the same spots in other stocks are inherited in that way.

Effect of temperature on the male spotting

During the course of the selection experiment it was fairly evident that some environmental factor was influencing the male spotting. Observations indicated that temperature was such a factor and to test this out a series of experiments were planned. These experiments were carried out at the tenth generation of selection when the divergence of the high and low strain was about as extreme as at any time during the selection experiment. The strains 500 low and high and 502 low and high were used in the tests. It had been noted that low rather than high temperature seemed to affect the results ordinarily obtained under room temperatures, so attention was directed chiefly to the effects of low temperatures as compared with room temperature.

A set of matings was made from each strain, some of them being kept under room temperatures (17°—23°C), and others being placed in a basement room where the temperatures ranged between eleven and

TABLE 12

Effects of temperature on spotting of males.

A. Strain 500 low

Temperature condition	Type 1	Type 2	Type 4	Type 5	Type 6	Type 11	Type 12	Type 13	Type 15	Type 16	Total males
Room (17-23° C)	546	5	2	6	8	9				3	579
Basement (11-15° C)	69					418	6		3	4	520

B. Strain 500 high

Room (17-23° C)	99	10	12	23	28	49	23	6	24	422	696
Basement (11-15° C)						54	10		15	73	152

C. Strain 502 low

Room (17-23° C)	92		1	4	1	166	7	1	4	8	284
Basement (11-15° C)	4					249			1		254

D. Strain 502 high

Room (17-23° C)									1	502	503
Basement (11-15° C)						15	8		5	97	125
Incubator (25° C)									1	61	62

In table 12 are shown the results of a series of experiments to test the effect of temperature on the spotting of males. Under room temperatures strain 500 low was almost pure for type 1 males, while strain 502 high was almost pure for type 16 males. Strain 500 high gave a fairly high percent of type 16 males under room temperatures while strain 502 low gave a good percent of type 1 males. From the tables it will be seen that the lower temperatures caused an increase in the type 11 males or in other words they brought out the outside pair of spots in all strains in which it did not already appear.

fifteen degrees centigrade. The species will not breed under temperatures much lower than those in the basement. The results are shown in table 12. No effects of high temperature had been observed in the laboratory, but to test the effect of high temperatures one culture of 502 high was reared in an incubator where the temperature was kept at 25°C. Strain 500 low which at room temperature produced almost exclusively males of the two-spotted type, reacted quite differently at lower temperatures. When 500 low was reared at the basement temperatures a large percentage of the males possessed very distinct outside spots. The middle pair of spots was not affected by the low temperatures. Strain 500 high which at room temperatures ordinarily produced a large

majority of six-spotted males, at basement temperature produced no two-spotted males at all, but only males having the outside pair of spots. Strain 502 low never produced as high a percentage of two-spotted males as the other low strains. This strain proved to be always more susceptible to the effect of low temperatures. The count for room temperature in table 12 is hardly characteristic of the strain, for at that time the percentage of two-spotted males was somewhat low. In the stocks of 502 low reared at the basement temperatures practically all of the males were those possessing the outside pair of spots. The stock of 502 high which at the time of the test was practically pure for six-spotted males when reared at room temperatures was also reared in the basement room. Since this stock was, at room temperatures, already pure for six-spotted males, low temperatures could not increase the percentage of males possessing the outside pair of spots. The results show that the low temperatures had no effect on the outside pair of spots but did seem to reduce somewhat the number of males possessing the middle spots. Instead of being pure for six-spotted males as was the condition under room temperatures a few of the males had one or both of the middle spots missing. Strain 502 high was also subjected to high temperatures. It was reared in an incubator where the temperature was kept at 25°C. Table 12 shows that results obtained at this temperature were not different from those obtained under room conditions.

These tests show that the outside pair of spots could be brought out by low temperatures regardless of the spotting condition of the strain induced by selection. Even in lines where none of the males under room temperatures possessed the outside spots, these spots could be made very distinct in a large majority of the males under lower temperatures. In strains where the males did not possess the middle pair of spots under room temperatures, the lower temperatures did not bring out these spots as it did the outside pair. Strains which under room temperatures possessed the middle pair of spots showed a tendency to lack them at low temperatures. In fact the lower temperatures showed a slight tendency to reduce the distinctness of this pair of spots. The lower temperatures also had a tendency to intensify the general pigmentation of the fly. Temperatures higher than room temperatures seemed to have no effect upon either the middle or outside pair of spots. It is true that the data on the effects of higher temperatures are rather meager and more matings would have given more conclusive results. It was the intention to test this out further at a later date but pressure of other work has made

this impossible. However, judging from observations made during the course of the selection experiment, I feel confident that more extensive data would only substantiate the results already recorded.

The facts brought out by this experiment have a definite bearing upon the selection experiments. Low temperatures occasionally encountered in the laboratory would have considerable influence upon the results of the selection experiment. This effect, especially upon the low strains, was noted several times. The effect of low temperatures is responsible for some of the irregularities in the graphs representing the behavior of the strains (see figures 2, 3 and 4). The rather sudden upward divergence of the graphs for the low strains near the end of the selection experiments can be correlated with a period of low temperatures which decreased the percentage of two-spotted males.

Crosses between the various male strains

At the end of the selection experiment when the various male strains had become fairly well established, crosses were made between some of them. A large series of experiments were planned but only a limited number were carried out because of the interference of other duties. Two types of matings were made, one set in which the high and low strains of the same line were crossed and the other in which the high strains of two different lines were used.

For the set of matings where the high and low strains of the same line were crossed, line 500 was used. Generation fifteen of the 500 low was crossed with generation fourteen of strain 500 high. The cross, 500 low male by 500 high female, and its reciprocal were both made and the results from the two types of crosses were the same (table 13). As to dominance or recessiveness, it would hardly be expected that either strain would act as a complete dominant for neither was absolutely pure for its type. The F_1 results show a high percentage of the males to be of the two-spotted type, but there are also considerably more six-spotted males than were to be found in the low strain. So the results show spotting in the males which is intermediate between the two strains. The lack of Mendelian ratios is probably due to the lack of dominance of either type. This would affect not only the F_1 generation but also the ratios obtained in the F_2 generation.

For the cross between two high strains, lines 500 and 502 were used. The fourteenth generation of the 500 high was crossed with the same generation of the strain 502 high. At the time of the cross the 502 high

TABLE 13

F_1 generation of crosses between various male strains.

A. Strain 500 high × strain 500 low

Type of cross	Type 1	Type 2	Type 4	Type 5	Type 6	Type 11	Type 12	Type 13	Type 15	Type 16	Total males
500 high ♀ × 500 low ♂	245	17	1	20	12	34	10		1	59	399
500 low ♀ × 500 high ♂	465	17	1	18	11	47	8		5	40	612
Total	710	34	2	38	23	81	18		6	99	1011

B. Strain 502 high × strain 500 high

502 high ♀ × 500 high ♂	5	2			1	14	5			425	452
500 high ♀ × 502 high ♂	4			1		3	1		1	136	146
Total	9	2		1	1	17	6		1	561	598

The results in table 13 are for the crosses between the high and low strains of line 500 and between the high strains of lines 500 and 502. The condition of the two strains of line 500 at the time of this cross is shown in the last generation of selection of these strains (tables 1 and 2). The cross gave a result which was intermediate between the two strains. The cross between the two high strains gave a result which might be considered like either strain since the two strains were very similar at the time of the cross.

strain was practically true for the six-spotted type of males while strain 500 high still threw a few two-spotted males in each generation. The cross was made in both directions with no difference resulting between the two types of crosses. The results found in table 13 show a very high percentage of six-spotted males. The results which are about characteristic of either strain show that the same factors are responsible for the spotting character in each strain.

Summary

1. Selection was effective in changing the percentage of two-spotted or six-spotted males in a strain. There was a difference in amenability to selection of the different lines taken from nature.

2. The effects of selection remained permanent, at least for several generations.

3. The crosses with the 501 type of male indicate that there is sex-linked inheritance involved in the case of both the middle and outside pairs of spots.

4. Low temperatures tend to bring out the outside pair of spots on the male and also seem to have a slight tendency to cause the middle pair of spots to disappear.

5. Neither type is dominant in the F_1 generation when strains with two-spotted males are crossed to strains with six-spotted ones.

6. The results of crosses between the high strains of different lines indicate that their spottedness is due to the same factors.

FEMALE SPOTTING

Abnormal female

While examining flies on March 16, 1916, in one bottle, six females were found which had the middle pair of spots missing from the fourth abdominal segment. The other two pairs of spots were normal in size and position but there was no indication of the middle pair (plate 1, no. 5, page 64). These females were mated to brothers and from this mating originated a strain of flies which was called the "abnormal female" strain or for the sake of brevity the "ab" strain. This strain is characterized by the absence of the middle pair of spots from the fifth abdominal segment.

Since there is sexual dimorphism in the spotting of the fifth abdominal segment, the problem of analyzing the spotting inheritance became rather difficult. In mating the original ab females to brothers, the males had to be selected blindly, as there was no way of knowing what relation if any, the male spotting bore to the female spotting. So in selection for purity with respect to the ab condition, the type of males used was kept constant, only those being used which possessed but one pair of spots, the large dorsal ones (plate 1, no. 1).

Selection of ab strain

As already mentioned the six original ab females were mated to two-spotted brothers in stock culture and the total of their offspring showed 576 normal females and 105 ab females. (It might be said here that these early counts of ab females are not absolutely accurate as there are always a few flies which have the ab spotting on one side and normal on the other. These intermediates at first were called ab or normals depending upon the intensity of the spots. Later they were recorded in the counts.) It was hoped that by mating ab females to their brothers a pure strain, with respect to the ab female, might be established. At first the results seemed rather puzzling for in each cross ab females were

TABLE 14

Selection from the original abnormal females to establish a pure abnormal strain.



The letter "N" in table 14 indicates normal females and "ab" stands for abnormal females. The figures above the line indicate the mating number in the generation. This selection line was lost at the end of the fifth generation.

obtained but the relative number of ab and normal females was quite variable. Table 14 shows the selection which was carried on for five generations. It will be seen that in the fourth generation of selection there were two matings (matings 5 and 18) which gave only ab females. Similar results were obtained in the fifth generation. Unfortunately at this time the selected strain was lost, due to the extreme summer heat. Previously, however, the ab strain had been crossed to a wild strain which did not show the ab condition in the females. From the F_1 of this cross a few ab females were saved and mated to brothers and in this way the selection was renewed.

It should be noted that the cross from which the ab strain was regained was one in which a male from the ab strain had been mated to a normal female from a wild strain (recorded as original cross in table 15).

Although the male himself did not show the ab character, he was capable of transmitting it to his offspring. In the F_2, or generation B, a large number of flies were obtained and the selection for a pure ab strain was once more started. In the third generation eighteen matings were made with the ab females as mothers, and in only one case (mating C 8) were the females obtained all ab and this was probably due to chance as the count was very small. In generation D the ab females were used in twenty-nine matings and six of them proved to be pure with respect to the ab condition. It was then thought that the strain had reached purity for the ab condition and thirteen matings were made from among the offspring of the matings which bred true. The results of this generation were again disappointing as the thirteen matings did not all breed true. Four matings were made from D27 and they were all pure, five were made from D44 and none of them were pure, four were made from D45 and two were pure. These inconsistencies could not be explained at this time but later matings added new light which made them explainable. From among the offspring of D27 more matings were made all of which bred true again. This is the origin of the ab strain which has bred true for many generations.

While the strain producing the ab females was being selected for purity, back selection was also attempted. This was to see if selection was effective in both directions. From mating B1 (table 15) back

TABLE 15

Selection in strain producing abnormal females.

Generation	Parent	Female spotting				Generation	Parent	Female spotting			
		Normal	Ab	Left	Right			Normal	Ab	Left	Right
Original cross		10	9			E7	D2	1	135	3	5
						E8	D15		130		1
A2		6	7			E11	D15	22	171	5	5
B1	A2	200	138	34	31	E13	D30		17		
C3	B1	51	92	18	11	E15	D31		117		
C4	B1	46	48	8	5	E16	D44	8	94	10	4
C5	B1	70	36	6	11	E17	D27		69		
C6	B1	16	65	9	10	E18	D27		124		
C7	B1	105	30	10	6	E20	D27		126		
C8	B1		5			E21	D27		128		
C10	B1	59	47	10	7	E23	D44	13	62	2	3
C12	B1	1	14	2		E24	D45	57	165	5	7
C13	B1	51	13	3	1	E27	D15	7	349	12	11
C16	B1	8	16	2	1	E28	D30		135		
C17	B1	55	9	8	2	E29	D30		125		

TABLE 15 (continued)

Generation	Parent	Female Spotting				Generation	Parent	Female spotting			
		Normal	Ab	Left	Right			Normal	Ab	Left	Right
C18	B1	54	20	5	4	E30	D30		115		2
C19	B1	24	33	7	5	E33	D33	1	179		1
C21	B1	2	11	2	2	E34	D33	4	153	2	
C22	B1	26	33	7	4	E36	D44	24	159	6	12
C26	B1	9	69	6	6	E37	D44	31	155	9	3
C27	B1	3	14	1		E39	D44	2	251	1	
C32	B1	2	18	1	1	E40	D45	1	237		1
D2	C1	8	81	4		E41	D45		198	1	
D3	C4	10	26	5	3	E42	D45		63		
D4	C4	39	67	3	9	E47	D4	27	242	8	7
D6	C4	29	46	6	3	F1	E18		161		
D7	C3	46	16	2	4	F2	E20		176		
D8	C3	17	44	4	7	F3	E20		84		
D9	C3	26	14	2	3	F4	E20		82		
D11	C6	2	16	4		F5	E20		175		
D12	C5	14	14	2	3	F6	E28		222		
D13	C5	25	30	4	2	F7	E28		76		
D15	C6	43	50	6	11	F8	E28		254		
D16	C7	3	41	2	2	F9	E28		178		
D17	C3	3	64	1	2	F11	E41	1	316	4	2
D18	C4	7	37	4	2	F13	E41	37	192	8	7
D22	C12		81	1		F16	E25	66	251	19	13
D26	C8	25	250	7	13	F17	E37	6	133	5	5
D27	C12		98			F18	E37	2	106		1
D30	C12	1	124			F19	E37	15	83	7	3
D31	C8	7	16	3		F25	E25	10	48	4	2
D33	C12	9	23	2	3	F26	E24		240	2	
D37	C6		27		1	F29	E24	84	159	22	16
D38	C6	18	36	3	3	F30	E41	15	207	12	3
D44	C26		253	1		F31	E29		414		1
D45	C26		138	2		F32	E21	2	370		
D46	C32		93			G1	F2		195	1	
D47	C32	10	31	2	1	G2	F2		176		
D49	C32	16	37	4	6	G3	F8		401		
D50	C26	9	145	1	1	G4	F2		66		
D51	C32	5	30	1	2	G5	F8		100		
E1	D15	10	285	8	12	G6	F2		128		
E2	D15	14	288	11	13	G7	F8		59		
E3	D15		247	1		H1	G3		186		
E4	D2		57			H2	G3		198		
E5	D2	1	4		2	I1	H1		67		
E6	D2	3	59	2		I2	H2		214		

Table 15 shows the selection required to bring the abnormal strain to purity. The letters in the column to the left indicate the generation of selection and the figures which follow them indicate the mating number in the generation. "Left" indicates a female which is abnormal only on the left side and "right" one which is abnormal on the right side.

TABLE 16
Back selection of abnormal strain.



Table 16 gives the results of selection for the normal female in the strain producing a high percent of abnormal females. The selection here is in a direction opposite to that in table 15 but started with mating B1 of that table. At generation 6 selection was again reversed (selecting for ab females) and is shown as the offspring of mating 4 of generation 6. The selection for normal females was still continued and is shown as the offspring of mating 6 of generation 6.

selection was started, using normal females. At the time back selection was started the strain was throwing about an equal number of ab females and normal females. Selection of this sort was carried on for nine generations and the strain became practically pure for normal females (table 16). In the fifth generation of back selection when the strain was approaching purity for the normal females, selection was started again in the reverse direction. This was done by using ab females as mothers. Selections in opposite directions were carried along for four generations, by which time one strain had become practically pure for the normal female, while the other was again throwing a large percentage of ab females. These experiments show that selection is effective in either direction before purity is reached and that purity may be reached for either type if selection is carried far enough.

The question as to how the ab strain originated,—whether as a simple mutation from the beginning, or as a result of selection—will not be discussed at present. Before taking up this question we will examine the results of the matings of this strain to normal strains. These matings were made in attempting to analyze this variation.

Matings of the ab strain with the normal strain 500 low

The strain 500 low to which the ab strain was crossed was one which had been bred in the laboratory for six generations and was in no way related to the ab strain. The males of this strain were almost all of what may be called the normal male type, having only the two large spots. The 500 low strain had been selected for three generations for the above type of male and from the beginning showed a tendency to produce males of that sort. The females of the 500 low strain were all normal, that is, having a full set of six spots. In all strains bred in the laboratory a few ab females appeared, but they seemed to be somatic variations as will be shown later (page 102). Since this strain was very similar to the ab strain, as far as the male spotting was concerned, and normal with respect to the female spotting, it was considered a good strain with which to cross in testing the genetic behavior of the ab strain.

As will be seen in table 17 twelve matings were made between abnormal females and males of the normal strain, and nine matings were made between males of the abnormal strain and females of the normal strain. For the first type of cross the totals were 977 normal females, 1645 ab females, 329 "left" females and 273 "right" females.[1] In the

[1] By "left" female is meant a female which is abnormal only on the left side and the expression "right" refers in the same manner to the right side.

TABLE 17

F_1 generation of cross abnormal strain × normal strain 500 low.

Abnormal ♀ × normal ♂					Normal ♀ × abnormal ♂				
Mating number	Female spotting				Mating number	Female spotting			
	Normal	Ab	Left	Right		Normal	Ab	Left	Right
150	57	158	17	17	157	164	190	42	46
151	64	81	30	19	158	204	206	56	39
153	15	142	19	15	159	35	91	21	21
154	72	124	28	21	168	66	177	27	32
155	79	187	25	29	169	123	329	51	59
162	24	103	14	13	170	75	261	43	42
163	57	77	17	11	171	24	158	13	10
164	53	92	22	24	172	52	183	30	22
165	91	26	13	8	250	154	230	60	55
166	92	139	33	26					
188	281	227	71	49					
251	92	289	40	41					
Totals	977	1645	329	273	Totals	897	1833	343	326

Grand total, 1874 normals, 3478 abnormals, 672 lefts, 599 rights.

In table 17 are shown the F_1 results of a cross between the abnormal female strain and strain 500 low. The females in strain 500 low were all normals. Crosses in both directions are shown. Both normal and abnormal females appeared in the F_1 generation with about twice as many abnormals as normals.

second set of matings in which the male from the ab strain was used the totals were 897 normals, 1833 abnormals; 343 lefts and 326 rights. These matings were all single-pair matings. The total of the F_1 is 1874 normals, 3478 abnormals, 672 lefts and 599 rights. These F_1 results threw very little light upon the nature of the abnormal variation. They do show however that either the male or the female of the ab strain will transmit the ab character even though the males do not show it. It is also seen that the ratio of abnormals and normals is about the same regardless of which way the cross was made.

In order to carry the analysis still further F_2 matings were made (table 18). Since there were both ab and normal females in the F_1 it was necessary to make two types of matings in the F_2, that is, ab females with brothers and normal females with brothers. The totals for the first type of matings were 945 normals, 1248 abnormals, 138 lefts and 112 rights, and for the second type 1879 normals, 1415 abnormals, 167 lefts and 159 rights. If the ab condition acts as a Mendelian character

there should be no difference in the genetic constitution of the two types of females in the F_1. Then F_1 ab and normal females when crossed should behave the same with respect to the sort of offspring to which they give rise. Three types of crosses were made in which the behavior of these two sorts of females can be compared: (a) F_1 females back-crossed to the pure ab strain, (b) F_1 females back-crossed to the normal strain which was used in the cross, and (c) F_1 females by brothers. The totals for the results of these crosses are shown in table 19. Considering the individual matings the results seem quite variable in each type of mating, but the totals show evidence of a slight difference in behavior of the two types of females appearing in the F_1. Where the two types of females were back-crossed to the pure ab strain there is a higher relative number of females, as compared with the number of normals produced when the ab female was the mother. Where the cross was a back-cross to the normal strain there was no apparent difference between the two females, but again when they were crossed to their brothers

TABLE 18

F_2 generation of cross abnormal female strain × strain 500 low.

Abnormal F_1 ♀ × normal brother					Normal F_1 ♀ × normal brother				
Mating number	Female spotting				Mating number	Female spotting			
	Normal	Ab	Left	Right		Normal	Ab	Left	Right
154 A1	80	125	20	13	154 A3	164	177	10	13
154 A2	73	177	5	9	157 A1	114	137	18	22
157 A3	167	157	27	20	157 A2	178	200	12	21
157 A4	126	101	15	8	157 A5	166	113	11	5
162 A1	107	179	15	16	157 A6	188	156	24	15
164 A3	63	172	14	11	164 A1	101	72	11	4
164 A4	105	213	19	16	164 A2	87	146	9	14
169 A1	83	28	5	5	169 A4	128	89	10	10
169 A2	141	96	18	14	188 A1	262	148	33	17
					250 A1	317	88	13	20
					251 A1	174	89	16	18
Totals	945	1248	138	112	Totals	1879	1415	167	159

Grand total, 2824 normals, 2663 abnormals, 305 lefts, 271 rights.

In table 18 showing the F_2 results of the cross abnormal female strain by strain 500 low, the grand total shows about an equal number of abnormal and normal females. However the ab F_1 female gave a higher percentage of abnormals in the F_2 than the normal female from the same generation.

a higher relative number of ab females appeared when the maternal parent was an ab female. With the large number of individuals examined in these crosses it hardly seems possible that these variations could be due to chance. Assuming that ab is an ordinary Mendelian character, these two types of F_1 females should react the same genetically, but when they are each crossed to males of the same type, they each tend to beget young which show a higher percentage like their mothers. Although there is a variation in their behavior the preponderance is never large enough to obliterate any class of individuals which should have

TABLE 19

A comparison of the behavior of the two types of females appearing in the F_1 generation.

Type of male used in cross	F_1 ab female				F_1 normal female			
	Female spotting				Female spotting			
	Normal	Ab	Left	Right	Normal	Ab	Left	Right
Male from pure normal strain	2706	382	142	138	2812	486	133	127
Male from pure ab strain	203	2922	110	81	223	1588	103	81
F_1 brother	945	1248	138	112	1879	1415	167	159

Table 19 shows the results of three types of crosses comparing the behavior of the two types of females found in the F_1 generation. In two of the crosses the F_1 ab female produced more abnormals than her normal sister.

segregated out. The factor which brings about the difference seems to play entirely upon the heterozygous individuals and therefore does not prevent the ab condition from still being considered as behaving in a Mendelian manner.

The F_2 normal and ab females were back-crossed to the respective strains from which they came, to test whether there was Mendelian segregation. These results are shown in table 20. Where the F_2 ab female was back-crossed to a male from the ab strain, eight of the fifteen matings were pure for the ab condition. Where the F_2 normal female was back-crossed to a male from the normal strain, nine out of the fifteen matings bred practically pure for the normal condition. The same sort of back-crosses were made in the F_1 generation (see table 19). Out of ten crosses between F_1 ab females and males from the ab strain no matings went pure for the ab condition and likewise no all-normal counts were obtained from making eight crosses between F_1 normal females and males from the normal strain.

TABLE 20

Back crosses of the two types of females appearing in the F_2 generation.

F_2 ab ♀ × pure ab ♂					F_2 normal ♀ × pure normal ♂				
Mating number	Female spotting				Mating number	Female spotting			
	Normal	Ab	Left	Right		Normal	Ab	Left	Right
205		141			210	398	10	10	14
206	19	109	2	3	211	56		4	2
207	22	94	9	7	212	321	10	7	15
208		157			213	387	5	6	4
209		39			214	220	21	9	10
220	14	222	10	4	218	208	57	18	12
221		67			225	369			
222	3	61		1	226	314		1	
223	17	157	9	7	227	227	1	1	3
224		223			228	269	49	18	15
241		86	1		229	210		1	
256	43	251	5	10	263	361	1		1
266	20	268	9	17	264	336		1	1
269		80			265	278	2	4	3
270		113			267	263		1	1

The crosses in table 20 were to test the segregation of the normal and abnormal females in the F_2 generation. Ab females from the F_2 generation were back-crossed to males of the pure ab strain while normal females of the same generation were back-crossed to males of the pure normal strain. Of the first cross, eight out of fifteen of the females acted as pure abnormals and in the latter cross, nine out of fifteen of the females acted as pure normals.

The fact that pure stocks were regained by back-crossing the two types of F_2 females to the stock from which they originated, indicates that there is Mendelian segregation. That this could not be done with apparently the same types of females from the F_1 also shows Mendelian behavior.

Some of the F_2 females having a spot on one side only (recorded as lefts or rights) were tested as to their genetic constitution. The test was made by back-crossing them to the original strains from which they came. The number of such crosses was not large but in all of them both abnormal and normal females appeared, showing that the females used were heterozygous flies. So the flies indicated in the tables as lefts and rights are probably individuals which are heterozygous for the ab and normal condition.

Summary of various types of matings in testing the behavior of ab when crossed to the strain 500 low

1. In the F_1 generation resulting from the cross between the ab female and the 500 low strain and its reciprocal, there is not a dominance of one or the other of the contrasted characters. Both types appear in F_1 with about twice as many abnormals as normals. Both males and females from the ab strain are capable of transmitting the ab character although only the females show it.

2. The two types of F_1 females when back-crossed to their respective original strains did not give pure stock, showing that these females were heterozygous and there was no segregation in this generation.

3. The two types of F_1 females gave slightly different ratios when crossed to males of the same genetic constitution. The F_1 ab female gave more abnormals than the F_1 normal female when they were crossed to the same male.

4. There was Mendelian segregation in F2 of the two original types. This was shown by back-crosses of the two types of females in F_2 and results of the F_2 generation.

Crosses between the ab strain and strain 502 high

In an attempt to further analyze the genetic behavior of ab, the ab strain was mated to strain 502 high. The strain 502 high was like 500 low so far as the female condition was concerned, that is, all the females were normal, i.e. with a full set of six spots. The only difference between these strains was that the males in strain 502 high had been selected for a full set of six spots while the strain 500 low had been selected for the normal or two-spotted male condition.

To the writer's surprise entirely different ratios were obtained (table 21) in F_1 when the ab strain was crossed to the strain 502 high. With apparently the same environmental conditions as before, a large majority of the F_1 flies were normal. There was practically a dominance of the normal condition over the ab. Out of the 1965 F_1 females examined only 71 were abnormal.

These F_1 results are quite different from those obtained when ab was crossed to strain 500 low (table 17). The count for the F_1 generation of the cross of ab with 500 low was 1874 normals, 3478 abnormals, 672 lefts and 599 rights. For the abnormal-by-502-high cross the counts were 1756 normals, 71 abnormals, 69 lefts and 69 rights. Since there is almost complete dominance of the normal condition over the abnormal condition in crossing ab by the 502 high strain, it would be expected that

TABLE 21

Totals of the F_1 and F_2 generations of the cross abnormal strain × strain 502 high.

F_1 generation			
Female spotting			
Normal	Ab	Left	Right
1756	71	69	69
F_2 generation			
3169	1117	54	47

The counts in table 21 are the results of the F_1 and F_2 generations of the cross between strain 502 high and the abnormal female strain. Very few abnormals appeared in the F_1 generation and an approximate 3:1 Mendelian ratio was obtained in the F_2 generation. Note that the ratios in this cross are quite different from those obtained in the cross, abnormal female by strain 500 low.

the usual 3 to 1 ratio would be obtained in the F_2. This was found to be true as will be seen in table 21. The totals for the F_2 were 3169 normals, 1117 abnormals, 54 lefts, and 47 rights. F_2 data were obtained from ab female by 502 high male and the reciprocal cross, but the ratios were approximately the same in each.

F_3 matings were also made to see whether a pure ab strain might be extracted. Fourteen matings were made of which two bred true to the ab condition. Since there is almost complete dominance of the normal condition, probably all of the ab females obtained in the F_2 are pure abnormals. There is no way to determine the genetic constitution of the males with respect to the ab condition except by breeding. One-fourth of the males should be pure for the ab condition, so one-fourth of the F_3 matings should have yielded only abnormals. Only two out of fourteen matings were of this character, but this shortage was probably due to chance because of the small number of matings.

Here we have the ab condition acting very much as an ordinary Mendelian character, the normal condition being almost completely dominant in the F_1 generation. The only thing unusual about this set of matings was that there was such a difference in behavior of the ab character here as compared with its behavior when crossed to strain 500 low. The explanation for this is found in the results of the tests which follow.

Genetic behavior of the ab strain when crossed to various strains

Since such a difference was found in the F_1 behavior of the ab strain when crossed to strains 500 low and 502 high, it was decided to test its behavior with all of the strains carried. The strains used were the selection strains of the experiment upon the male spotting. The females in all these strains were normal, having a full set of six spots. The only difference was that part of the strains had been selected for males with two spots while the rest were selected for the six-spotted condition of the males. When these matings were made the high strains had attained a condition in which a large percentage of the males were of the desired type with six spots, and likewise the low strains possessed mostly two-spotted males. The matings were all made at once and subjected to as nearly identical environmental conditions as possible. When the F_1 counts were made the same difference was again found to exist between the ratios when ab was crossed to 500 low and 502 high. Some other rather striking results were obtained from this series of crosses.

The results show that the dominance or recessiveness in the F_1 is affected by the degree of spottedness of the males in the strain in which the ab strain is crossed (see table 22). All three of the strains which are called high give almost complete dominance of the normal female condition over the ab in F_1 while the low strains crossed to the ab strain yield nearly half or more of abnormal F_1 females. The ratios obtained were as follows: In line 500 the low strain produced 307 normals to 289 abnormals, and the high strain 710 normals to 44 abnormals; in line 502 the low strain, 176 normals to 259 abnormals, and the high strain, 457 normals to 7 abnormals; in line 504 the low strain, 166 normals to 427 abnormals, and the high strain, 296 normals to 93 abnormals. Since this consistent difference appeared in the reaction of the high and

TABLE 22

Results of F_1 generation in crossing abnormal strain with male-selection strains.

Strain crossed with ab	Female spotting				Strain crossed with ab	Female spotting			
	Normal	Ab	Left	Right		Normal	Ab	Left	Right
500 low	370	289	60	72	500 high	710	44	28	18
502 low	176	259	23	24	502 high	457	7	12	8
504 low	166	427	37	45	504 high	296	93	20	15

Table 22 shows the difference in reaction of the abnormal female strain in the F_1 generation when crossed to the various strains of the male selection lines. The percent of ab females is much higher in crosses to the low strains than in the crosses to the high strains.

low strains in crosses to the ab strain when all environmental factors were kept constant, it seems that the difference in genetic constitution of the two strains is the responsible factor.

Relation of temperature to genetic behavior of abnormal female condition

After the ab strain became once established it was quite evident that variations of temperature had no effect upon it, for although there was considerable fluctuation in the temperature of the laboratory the strain remained pure for the ab condition.

The variability of the recessiveness or dominance of the ab in the F_1 led the writer to suspect that some environmental condition might be playing a part. In addition to the influence of the "spottedness" of the males in the strain to which it was crossed there seemed to be some other factor. When the type of male to which the ab was crossed was kept constant there was still considerable variation in the F_1 results. This was most noticeable in crosses where strains possessing normal or two-spotted males were used, resulting in F_1 ratios with a high percentage of ab females. Here were obtained rather variable results. Temperature being one of the simplest environmental factors to analyze, a series of experiments were carried out to test its influence upon the behavior of ab in the F_1 generation.

In testing the effect of temperature, flies reared at room temperature were compared with some reared at high temperature in an incubator, and also with others reared at a low temperature in a basement room without heat. The effects of temperature upon the F_1 behavior of ab when crossed to strains 500 low and 502 high were tested. The results of these two experiments are given in table 23. The only place where it was possible to keep the temperature constant was in the incubator. The laboratory and basement temperatures were somewhat variable but the range of variation was sufficiently different to make a satisfactory experiment. The incubator was kept at 25°C which was just about the maximum temperature at which it was possible to rear *Drosophila busckii*. Even at that temperature the flies were somewhat undersized and reduced in number. Immediately after pairing the flies, the culture bottle was placed in the incubator and kept there continually except while the progeny was being examined. In the case of the cold, the flies were also reared at about the minimum temperature at which they would reproduce. The basement-room temperatures were so low that it was difficult to get the flies to mate when kept there from the beginning. To overcome this the flies

TABLE 23

Effect of temperature on the F_1 behavior of the abnormal strain.

F_1 results of cross ab × 500 low					F_1 results of cross ab × 502 high				
Temperature condition	Female spotting				Temperature condition	Female spotting			
	Normal	Ab	Left	Right		Normal	Ab	Left	Right
Room 17-23° C	370	289	60	72	Room 17-23° C	1299	64	57	61
Incubator 25° C	466	189	71	66	Incubator 25° C	142	1	2	4
Cold 11-16° C	3	625	8	4	Cold 9-15° C	265	133	23	42

Table 23 shows the effect of temperature on the F_1 behavior of the abnormal strain when crossed to strains 500 low and 502 high. It has already been shown that the ab strain reacts differently with each of these strains in the F_1 generation under normal room temperatures. From the table it is seen that lower temperatures increase the percent of ab females in the F_1 generation in each of the crosses.

were started in the laboratory and kept there till several eggs were laid. In the cross between 500 low and ab, in the low-temperature series, the culture bottles were kept at laboratory temperatures for the first week and then kept in the cold for the rest of the time. At that time the basement temperature varied between eleven and sixteen degrees centigrade. When the experiment was carried out with the strain 502 high the basement temperatures were somewhat lower (9°—15°C) and it was necessary not only to start the cultures in the laboratory but also to alternate cultures between the laboratory and cold room.

The results obtained from these matings are as follows: High temperatures tend to increase the number of normal females in the F_1 generation while low temperatures tend to increase the number of abnormals. As will be seen in table 23 giving these results, the temperatures seem to have a greater or less range of effect. Under room temperatures, strains 502 high and 500 low, due to a difference in spottedness, reacted differently in the F_1 generation when crossed with ab. Under the different temperature conditions they still retained their characteristic influence upon the F_1 behavior of ab. The lowest temperature used made the ab condition almost completely dominant when crossed to 500 low while the highest temperatures used made the ab condition almost completely recessive when crossed to 502 high. This is a very pretty demonstration of the combined influence of two factors upon the behavior of a Mendelian character.

Since temperature influences so much the ratios of the F_1 generation it is evident that its influence is upon heterozygous flies. To further demonstrate this some bottles of the F_2 generation of the cross 502 high by ab were reared in the cold. In this case a larger percentage of the females should be of the ab type than when the F_1 generation of the same cross was reared under the same condition. In the F_2 generation some pure abnormals should segregate out and not being influenced by the low temperature would increase this class. The results of the F_2 generation reared under the low temperatures were as follows: 398 normals, 398 abnormals, 18 lefts and 17 rights. The results of the F_1 generation reared under the same conditions were (table 23); 265 normals, 133 abnormals, 23 lefts and 42 rights. It is seen that the expected increase in the percentage of abnormals in the F_2 generation occurred. This is additional evidence that temperature in modifying the ratios obtained from crosses of the ab female strain, acts upon the heterozygous flies.

Linkage

Since the writer (WARREN 1917) had earlier found two eye mutations in this species it was thought that crosses between these two mutations and ab might be of interest. Earlier crosses between these two eye mutations indicated that they were probably in the same chromosome. The two eye mutations had been called "red" and "chocolate."

Ab was first crossed to the mutation red eye. In the F_1, as had been observed in other crosses, the ab was only partially recessive to the normal spotting in the red-eye strain. In the F_2, in case the red-eye mutation and ab were in the same chromosome we would expect to get no red ab females (the males cannot be considered since they do not show the ab character). If the two mutations should not be in the same chromosome we would expect independent segregation of the two characters in the F_2 and the chances would be that about one-half of the red-eyed flies would be abnormals. The F_2 results (table 24) were as follows: 193 red eye, normal-spotted; 110 red eye, ab-spotted; 640 normal eye, normal-spotted; 472 normal eye, ab-spotted. So it is seen that there is no linkage between red eye and ab, or in other words they are not in the same chromosome.

Ab was also crossed to chocolate eye. Since the red- and chocolate-eye mutations had been shown to be probably in the same chromosome we would expect ab to behave in the same manner when crossed to chocolate as it did with red eye. This expectation was realized, for chocolate and

TABLE 24

Crosses testing linkage between the abnormal character and two eye mutations.

A. F_2 of red-eyed strain × abnormal strain

Red-eyed females				Normal-eyed females			
Normal	Ab	Left	Right	Normal	Ab	Left	Right
193	110	12	12	640	472	47	34

B. F_2 of chocolate-eyed strain × abnormal strain

Chocolate-eyed females				Normal-eyed females			
Normal	Ab	Left	Right	Normal	Ab	Left	Right
102	121	14	10	319	398	37	47

Table 24 gives the results of crosses to test the linkage between the abnormal character and two eye mutations. Since there was independent segregation of the two characters in both crosses it can be assumed that the factor for the ab character is not in the same chromosome as the two eye mutations.

ab segregated independently in the F_2 generation. The results were as follows: 102 chocolate eye, normal-spotted; 121 chocolate eye, ab-spotted; 319 normal eye, normal-spotted; 398 normal eye, ab-spotted. So the indications from these crosses are that the mutation for the ab condition was not in the same chromosome in which the red and chocolate eye occurred. Since no other mutations have been found in this species it was impossible to test the linkage further.

Attempts to establish an ab strain from other stocks

In practically every strain of flies which was carried any length of time there appeared an occasional female which had the ab type of spotting. In some cases, out of a day's count of sixty females from a bottle there might be four or five females with the ab type of spotting. So high a percentage was unusual, for sometimes several hundred females could be examined without a single female of that type appearing. However the relative number of the ab type of females appearing was a factor of considerable fluctuation and seemed to be influenced by some environmental condition. The environmental condition necessary to bring them out was not determined although close observation was made with this point in mind.

During the course of this experiment, as these females of the ab type

appeared in the stocks from time to time, some of them were isolated and mated in an attempt to establish another ab strain. In some cases these strains were selected for several generations. Also some of the females of the ab type were crossed to males from the pure ab strain.

The data for these attempts are given in table 25. The strain 508 was selected for seven generations for the ab condition, but at the end of this period the strain was no nearer purity than at the beginning. In the sixth generation seven single-pair matings were made, but none of these

TABLE 25

Attempts to establish by selection a second abnormal strain.

A. Selection for ab in strain 508

Generation and mating	Female spotting				Generation and mating	Female spotting			
	Normal	Ab	Left	Right		Normal	Ab	Left	Right
Gen. A mating 1	124	14	3	4	Gen. F mating 2	163	11	7	12
Gen. B mating 1	219	17	13	14	Gen. F mating 3	106	9	9	10
Gen. C mating 1	217	7	6	9	Gen. F mating 4	39	9	10	4
Gen. D mating 1	252	28	24	18	Gen. F mating 5	224	24	27	23
Gen. E mating 1	160	44	22	25	Gen. F mating 6	57	6	5	10
Gen. F mating 1	61	4	6	7	Gen. F mating 7	80	10	12	14

B. Selection for ab in strain 503

Generation	Normal	Ab	Left	Right
A	289	4	3	3
B	223	11	5	3
C	147	1		1

Table 25 shows two attempts to establish a second ab strain from occasional ab females appearing in strains 508 and 503. Although strain 508 threw an unusual number of ab-appearing females, this percent could not be increased in six generations of selection. In strain 503 only a few abnormals ever appeared.

matings showed any high percentage of abnormals. Again in strain 503, selection was carried on for three generations with no effective results. In three other cases ab females from the normal strains were mated to brothers but among the offspring there were very few if any abnormals

Another type of mating was made to test the genetic constitution of the ab type of females appearing in the normal strains. This was to cross the ab-appearing females to males from the pure ab strain. If these females were of the same genetic constitution as the females in the pure ab strain, a cross to the males in the ab strain should breed true for the ab condition. The results of these crosses (table 26) were, that no more abnormals appeared in the F_1 than ordinarily appeared when the males from the ab strain were crossed to normal females from the respective strains in which the ab females were found. So it seems that

TABLE 26

Crosses of abnormal-appearing females found in wild stock to males from the pure abnormal stock.

Types of female spotting			
Normal	Ab	Left	Right
109	95	20	28
60	149	19	25
67	327	23	24
68	194	16	18

The counts in table 26 give the F_1 results of crosses of the occasionally appearing ab females in wild stocks to males from the pure abnormal stock. Had these ab-appearing females been genetically pure abnormals all the offspring of the F_1 generation should have been abnormals. No higher percent of abnormals appeared in the F_1 generation than if the mating had been made to normal-spotted females from a wild strain.

these constantly appearing females of the ab type in the various strains were not genotypically abnormals.

Discussion of results of the analysis of the abnormal type of female spotting

From the very beginning there existed two peculiarities of the abnormal strain which made the analysis of the character both difficult and confusing. These were sexual dimorphism in the spotting of the species and the variability of the dominance or recessiveness in the F_1 generation when crossed to a normal strain.

Discussing first the origin of the ab strain,—a glance at the data might seem to indicate that the purity of the ab strain had been arrived at by the cumulative effect of selection. The data shows that it required several generations of selection to bring the strain up to a state of purity and also that a somewhat gradual change took place in the direction of

selection. However it seems to the writer that the two peculiarities mentioned above offer an adequate explanation for the selection required to bring the ab strain to purity. Since the males did not show the character being selected for in the females, selection was necessarily blind as far as the paternal parent in each generation was concerned. This, together with the fact that heterozygous females might have the same appearance as pure ab females, would furnish sufficient explanation for the period of selection required to bring the ab strain to purity. The lack of success in attempting to establish pure ab strains from the occasional ab type of female appearing in normal strains indicates that the original ab individuals were probably mutations and the later-appearing were only somatic variations brought about by some environmental condition.

As to the genetic behavior of ab, early matings seemed to indicate that it had but few things in common with the behavior of a single Mendelian character. This was due largely to the lack of complete dominance or recessiveness of the ab in the F_1 generation when crossed to normal. More confusion was added when it was found that the relative number of abnormals appearing in the F_1 was influenced by the type of male spotting in the strain to which it was crossed. But after all of the data were summarized testing the effect of temperature and male spotting the following findings seem to hold true: *The ab condition in the female behaves as a Mendelian character. Its dominance or recessiveness is affected by temperature and by the type of male spotting in the normal strain to which it is crossed.* The effect of the spottedness of the males in the strain to which ab is crossed, is that the ab condition is almost entirely recessive in the F_1 generation when crossed to a normal strain in which the males are practically all of the type with a full set of six spots. The opposite holds true when ab is crossed to a normal strain where the males are of the two-spotted type. In this case ab acts more like a dominant over the normal female. High temperatures tend to reduce the number of abnormals appearing in the F_1 generation, while low temperatures increase them, but taking the effect of these factors into account the behavior of ab seemed to be that of an ordinary Mendelian character.

Summary

1. The type of female possessing the abnormal spotting probably originated as a mutation.

2. Although selection was necessary to bring the strain to purity for the abnormal type of spotting, the resulting purity was probably due only to isolation of individuals possessing the desired type of spotting, not to a cumulative effect on the strain as might be interpreted from the data. The difficulty encountered in bringing the strain to purity was due to the fact that the character was sex-limited.

3. The abnormal-spotting character acted as a Mendelian character. However the dominance or recessiveness of this character in the F_1 generation was rather variable and was found to be influenced by at least two factors.

4. One factor which influenced the behavior of the abnormal-spotting character in the F_1 generation was the strain to which the abnormal-female strain was crossed. When the strain possessing females with abnormal spotting was crossed to a strain possessing males of the six-spotted type, the former tended to act as a recessive character, but when crossed to a strain with males of the two-spotted type, it acted more like a dominant.

5. Temperature also influenced the behavior of the abnormal-spotting character in the F_1 generation. High temperatures tended to make the abnormal spotting act more like a recessive character while low temperatures cause it to act like a dominant one.

GENERAL DISCUSSION

The simplicity of a character offers no clue as to the simplicity of the manner of its inheritance. To the writer the most striking fact brought out in this study was the complexity of inheritance. A study of the spotting inheritance of a single body segment which bears only six spots might not ordinarily be expected to be very complicated. Numerous influences and interrelations of both genetic and environmental factors were found, but when carefully analyzed the inheritance was found to be Mendelian.

One of the most striking peculiarities was the almost entire independence of inheritance in the two sexes. Variations in the males of a strain did not usually influence the female spottng. The ab female and the 501 type of male were identical as far as the type of spotting was concerned. Each had the middle pair of spots missing but still there appeared to be no relation between the two when their inheritance was considered. Both males and females might possess the three pairs of spots but a different set of factors seemed to be responsible for the inheritance of the spots in each sex.

In all of the crosses between the various types of spotting studied in this paper there were no cases of complete dominance. The lack of dominance in the crossing of contrasted characters is frequently observed but its occurrence in all of the crosses in this work is worthy of note. In the case of the cross between the ab female strain and a strain of normal females, the percentage of ab females appearing in F_1 generation was dependent upon the temperature and the type of spotting of the male in the normal strain. Under certain combinations of these influencing conditions the ab condition was almost completely dominant.

Again each pair of spots seemed to be inherited independently and was acted upon differently by the same environmental factor. Low temperatures tended to intensify the outside pair of spots on the male while the same temperature tended to cause the middle pair of spots to be less distinct. Even the same pair of spots in the male was inherited differently in different strains. As simple a character as a pair of spots was dependent upon several factors.

The fact that one cannot look entirely to the germ cells for the solution of a problem in inheritance is rather strikingly brought out in this study. The importance of environmental conditions for the realization of certain characters was repeatedly seen. Temperature was shown to play an important part in making the ab condition a dominant or recessive character. The spotting in the males which seemed to be a character independent from the female spotting also influenced the dominance or recessiveness of the ab character. Again in the male spotting, certain spots which had been made to disappear by selection could be brought out again by subjection to certain temperatures. So when we consider the interrelation of different characters and the influences of environmental factors upon Mendelian characters, it is not surprising that the solution of the problem of inheritance has become more difficult than was anticipated at the time of the rediscovery of Mendel's Law.

In recent years considerable study has been devoted to the heredity of secondary sexual characters. Since the variations discussed in this paper are limited to one sex they may be considered as secondary sexual characters. It was found that the males were able to transmit variations in the female spotting which they themselves did not show. Likewise the females transmitted male variations which did not show. This ability of one sex to transmit the secondary sexual characters of the other has already been described in other insects. A similar case is that described by FOOT and STROBELL (1913) in the bug, Eu-

schistus. They crossed two species, one of which possessed a black spot on the male. The females of both species lacked the spot. However, the female of the species of the black-spotted male transmitted the spot although she herself did not show it. Other similar cases are GEROULD'S (1911) work on *Colias philodice* and DE MEIJERE'S (1910) work on *Papilio memnon*.

Considering the findings of this study in their relation to the evolution of the species it is difficult to say just what is happening. The spotting of the female seemed to be the more stable type. In the female the full set of six spots seemed to be the normal type while in the male one pair of dorsal spots was the commoner type in nature. The frequent variations found in the male spotting would indicate that evolution was taking place in that sex. In case an evolutionary process is in progress there would be two possibilities. The process might be one of loss or acquisition of spots to the fly. In the case of the species losing spots, all of the males would have once possessed six spots but have been the first of the two sexes to be acted upon by the process. On the other hand if the species be acquiring spots the action of the process upon the male has been retarded, the female having already changed from a two-spotted type to a six-spotted one. These are only suggestions of what might possibly be taking place, as there is no evidence from which we can draw any dependable conclusions.

GENERAL SUMMARY

1. This paper is a discussion of spotting inheritance in the fruit fly, *Drosophila busckii*. The spotting of the fifth abdominal segment is the subject of discussion.

2. There is sexual dimorphism in the spotting of this segment, the normal female having three pairs of spots on this segment and the male having one, two, or three pairs of spots on it.

3. An unusual type of female was found (probably originated as a mutant in the writer's stocks) which had one of the three pairs of spots missing. This was found to bear no relation to a similar type of spotting in the male; in fact the various types of spotting found in the two sexes were inherited independently.

4. The mutant in female spotting required considerable selection to bring the strain to purity, but the effect of selection was probably only to isolate individuals homozygous for the character.

5. Extensive selections were made for the type of male having one

pair of spots and the type having three pairs of spots. The strains in which selection was made for the two types of males were of a common parentage. By selection strains were produced which had a high percentage of the desired type but neither was ever brought to absolute purity for the type sought.

6. The behavior of the mutant in female spotting in the F_1 generation was affected by the strain to which it was crossed. If crossed to a strain possessing a high percentage of six-spotted males, it acted as a recessive character but if crossed to a strain with a high percentage of two-spotted males, it acted more like a dominant.

7. High temperatures caused the mutated type of female spotting to act as a recessive character while low temperatures made it act as a dominant.

8. In the male spotting, high temperatures intensified one pair of spots while the same temperatures seemed to reduce another pair of spots on the same segment.

9. In the male spotting, when a strain selected for two-spotted males was crossed to a strain selected for six-spotted males neither type of spotting was dominant in the F_1 generation.

10. There was a difference in amenability to selection of different strains taken from nature; but when two strains, each selected for six-spotted males but of different origin, were crossed, the results of the cross indicated that their spottedness was due to the same factors.

11. An unusual type of male spotting was found in which the spots of one pair were very large. A study of this character indicated that sex-linked inheritance was involved in its genetic behavior.

Acknowledgments are due Professor FERNANDUS PAYNE for many helpful suggestions and criticisms during the course of this study. Also I am indebted to my wife, ELMIRA SHIERLING WARREN, for much assistace both in the breeding work and preparation of material for this paper.

LITERATURE CITED

AGAR, W. E., 1913 Transmission of environmental effects from parent to offspring in *Simocephalus vetulus*. Phil. Trans. Roy. Soc. London B **203**: 319-351.

BRIDGES, C. B., 1916 Non-disjunction as proof of the chromosome theory of heredity. Genetics **1**: 1-52, 107-163.

CASTLE, W. E., 1910 The effect of selection upon Mendelian characters manifested in one sex only. Jour. Exp. Zoöl. **8**: 185-192.

1912 The inconstancy of unit characters. Amer. Nat. **46**: 352-362.

1916 Can selection cause genetic change? Amer. Nat. **50**: 248-256.

1917 a Piebald rats and multiple factors. Amer. Nat. **51**: 102-114.

1917 b The rôle of selection in evolution. Jour. Wash. Acad. Sci. **7**: 369-387.

DEXTER, J. S., 1914 The analysis of a case of continuous variation in Drosophila by a study of its linkage relations. Amer. Nat. **48**: 712-758.

DONCASTER, L., 1914 The determination of sex. pp. xi + 172. Cambridge: Cambridge University Press; New York: G. P. Putnam's Sons.

FOOTE, K., and STROBELL, E. C., 1913 Preliminary note on the results of crossing two Hemipterous species. Biol. Bull. **24**: 187-204.

GEROULD, J. H., 1911 The inheritance of polymorphism and sex in *Colias philodice*. Amer. Nat. **45**: 257-283.

HOGE, M. A., 1915 The influence of temperature on the development of a Mendelian character. Jour. Exp. Zoöl. **18**: 241-297.

JOHANNSEN, W., 1903 Ueber Erblichkeit in Populationen und in reinen Linien. 68 pp. Jena: Gustav Fischer.

1911 a Elemente der exacten Erblichkeitslehre. Zweite Ausgabe. 723 pp. Jena: Gustav Fischer.

1911 b The genotype conception of heredity. Amer. Nat. **45**: 129-174.

LUTZ, F. E., 1915 Experiments with *Drosophila ampelophila* concerning natural selection. Bull. Amer. Mus. Nat. Hist. **35**: 605-624.

MACDOWELL, E. C., 1915 Bristle inheritance in Drosophila. 1. Jour. Exp. Zoöl. **19**: 61-97.

1917 Bristle inheritance in Drosophila. 2. Jour. Exp. Zoöl. **23**: 109-146.

DE MEIJERE, J. C. H. 1910 Ueber Jacobsons Züchtungsversuche bezüglich des Polymorphismus von *Papilio Memnon*. Zeitschr. f. indukt. Abst. u. Vererb. **3**: 161-181.

MOENKHAUS, W. J., 1911 The effects of inbreeding and selection on fertility, vigor, and sex-ratio in *Drosophila ampelophila*. Jour. Morph. **22**: 123-154.

MORGAN, T. H., 1914 Heredity and sex. pp. viii + 284. New York: Columbia University Press.

1915 The rôle of environment in the realization of a sex-linked Mendelian character in Drosophila. Amer. Nat. **49**: 385-429.

1916 A critique of the theory of evolution. 197 pp. Princeton: Princeton University Press.

1917 An examination of the so-called process of contamination of the genes. Anat. Rec. **11**: 503-504.

MORGAN, T. H., STURTEVANT, A. H., MULLER, H. J., and BRIDGES, C. B., 1915 The mechanism of Mendelian heredity. pp. xiii + 262. New York: Henry Holt & Co.

MULLER, H. J., 1914 The bearing of selection experiments of Castle and Phillips on the variability of genes. Amer. Nat. **48**: 567-576.

PAYNE, FERNANDUS, 1918 An experiment to test the nature of the variations on which selection acts. Indiana Univ. Studies. Vol. 5, study 36.

SHULL, G. H., 1911 The genotypes of maize. Amer. Nat. **45**: 234-252.

WARREN, D. C., 1917 Mutations in *Drosophila buscckii* Coq. Amer. Nat. **51**: 699-703.

1918 The effect of selection upon the sex-ratio in *Drosophila ampelophila*. Biol. Bull. **34**: 351-372.

ZELENY, CHAS., and MATTOON, E. W., 1915 The effect of selection upon the "bar-eyed" mutant of Drosophila. Jour. Exp. Zoöl. **19**: 515-529.